WATER QUALITY DYNAMICS AND MANAGEMENT IN RESERVOIRS: A REVIEW OF RECENT MONITORING AND ADAPTIVE STRATEGIES

Author: **Rozibayeva Sadoqat A'zam qizi**
E-mail: ruzibayevas2026@gmail.com

Abstract

Reservoir water quality is controlled by interactions among watershed nutrient inputs, hydrological variability, thermal stratification, internal nutrient cycling and reservoir operations. This narrative review synthesizes 12 peer-reviewed publications from 2022–2025 to examine temporal and vertical water-quality dynamics and evaluate monitoring and management options. Evidence from field observations, process-based simulations, satellite remote sensing and machine learning indicates that surface observations alone cannot reliably characterize deep-water oxygen depletion or sediment-associated chemical changes. Sentinel-2 and Landsat products improve spatial coverage of optically active indicators, whereas profiling sensors and laboratory measurements remain essential for dissolved oxygen, nutrients and metals. Process-based models support scenario assessment, while data-driven methods are useful for interpolation but may be less reliable under unprecedented warming. An integrated, risk-based monitoring and adaptive management framework is proposed, combining watershed load reduction, vertical profiling, satellite surveillance and evaluation of selective withdrawal or hypolimnetic oxygenation. The review identifies the need for reservoir-specific calibration, uncertainty reporting and coordinated management of water quantity and quality.

Keywords: reservoir; water quality; thermal stratification; eutrophication; dissolved oxygen; remote sensing; adaptive management.

1. Introduction

Reservoirs provide drinking water, irrigation storage, hydropower and flood regulation, but these functions depend on water quality that can vary sharply by season, depth and location. Nutrient enrichment, suspended sediments, organic matter and temperature shifts interact with storage fluctuations and operational withdrawals. Reservoir-specific assessments are therefore necessary rather than assuming uniform water-column conditions [1–3].

Recent studies document persistent eutrophication and oxygen stress in potable-water reservoirs, and show that conventional periodic grab samples can miss short-lived or deep-water deterioration [1,4]. At the same time, advances in satellite observations, automated sensors and numerical modelling enable monitoring at complementary spatial and temporal scales [5–8].

The objective of this review is to (I) synthesize key mechanisms driving reservoir water-quality dynamics; (II) compare monitoring and prediction approaches; and (III) develop an integrated, practical management framework grounded in recent research.

2. Materials and Methods

A structured narrative review was conducted using 12 peer-reviewed journal articles published in 2022–2025. Publications were identified through targeted searches of journal and publisher records using combinations of "reservoir water quality", "stratification", "dissolved oxygen", "remote sensing", "machine learning", "selective withdrawal" and "oxygenation". Studies were included when they directly examined reservoir processes, monitoring, prediction or management; selected broader inland-water reviews were retained for methodological context. This was not an exhaustive systematic review or meta-analysis.

© 2026 London International Monthly Conference on Multidisciplinary Research and Innovation. All Rights Reserved.

The evidence was organized into four domains: (1) hydrological and thermal drivers; (2) nutrient–oxygen–sediment interactions; (3) observation and modelling methods; and (4) operational and catchment management. Findings were synthesized qualitatively, with numerical values reported only when explicitly presented in the cited studies. No new field samples, laboratory experiments or reservoir-specific model runs were performed.

Table 1. Key variables and their management relevance.

Indicator	Main process represented	Preferred measurement
Temperature profile	Mixing and stratification	Depth-resolved logger
Dissolved oxygen (DO)	Hypoxia and redox conditions	Profiling sonde / continuous sensor
Chlorophyll-a	Phytoplankton biomass	Laboratory + calibrated satellite
Total N / total P	External and internal nutrient supply	Laboratory sampling
Turbidity / TSS	Sediment transport and runoff	In situ + satellite
Fe / Mn	Redox-sensitive metal mobilization	Laboratory / spectral sensors

The proposed monitoring design distinguishes surface-visible optical signals from variables requiring direct measurements. It also considers seasonal sampling bias, satellite atmospheric correction, model extrapolation risk and the need for validation with independent observations [5–9].

2.1. Conceptual synthesis

The conceptual model (Figure 1) links upstream loading and reservoir hydrodynamics to measurable outcomes and management responses. It is a synthesis diagram, not a calibrated causal model.




Figure 1. Conceptual pathway from catchment pressures to reservoir water-quality management (author-developed synthesis).

3. Results

3.1. Seasonal and vertical dynamics

Across the reviewed evidence, seasonal thermal stratification emerges as a central control on water-quality variability. During stratified periods, surface and bottom waters may become physically isolated, allowing oxygen consumption to exceed replenishment in the hypolimnion. In a study of Yamchi Reservoir, summer bottom-water DO fell below 2 mg/L, while autumn and winter profiles were more uniform [2]. This result illustrates a site-specific mechanism rather than a universal reservoir threshold.

© 2026 London International Monthly Conference on Multidisciplinary Research and Innovation. All Rights Reserved.

Long-term monitoring of potable-water reservoirs further shows that nutrient enrichment, chlorophyll-a, organic carbon and diel DO dynamics can reveal deterioration not apparent from occasional samples [1]. High-frequency observations also detect rapid variation in iron and manganese related to oxygenation and seasonal redox changes [3]. These findings favor depth-resolved and high-frequency measurements during warm and transitional seasons.

3.2. Hydrology, climate and operating conditions

Water level, inflow timing and withdrawal depth can alter the thickness and temperature of deep-water layers, with consequences for oxygen availability. Modelling in Rappbode Reservoir showed that water-level reduction could aggravate hypolimnetic hypoxia and that selective withdrawal performance depends on timing and depth [4]. Comparative modelling indicates that machine-learning methods can reconstruct observed temperature profiles effectively, yet process-based simulations may be more suitable for extrapolation to novel warming scenarios [6].

3.3. Monitoring technologies

Satellite imagery provides repeated spatial observations of chlorophyll-a, suspended matter, turbidity and related optical indicators, although retrieval accuracy varies by sensor, atmospheric conditions and reservoir characteristics [5,7,8]. In a multi-reservoir study, random forest performed favorably against several alternative algorithms for chlorophyll-a, Secchi depth and suspended solids [7]. A separate study demonstrated that remote sensing combined with profile-shape information and machine learning can estimate vertical variables, reporting a test-set mean absolute error of 0.25 mg/L for DO in Wadi Dayqah Reservoir [9]. Such performance is study-specific and does not eliminate the need for in situ validation.

Table 2. Complementary monitoring approaches.

Approach	Strength	Principal limitation
Grab samples / laboratory	Direct nutrient and metal analysis	Sparse time coverage
Continuous depth sensors	Short events and vertical gradients	Maintenance and fouling
Satellite optical retrieval	Reservoir-wide spatial patterns	Surface bias; clouds; calibration
Machine learning	Nonlinear prediction and fusion	Transferability and extrapolation
Process-based model	Scenario testing and mechanisms	Input data and calibration demands

4. Discussion

4.1. Linking monitoring to management

The synthesis suggests that effective reservoir management should begin with a conceptual source–pathway–receptor model and an explicit decision context: drinking-water abstraction, ecological protection, irrigation suitability or multipurpose operation. For nutrient-dominated systems, reducing external nitrogen and phosphorus loading is a first-line intervention; otherwise, in-reservoir treatments may deliver only temporary improvements. For oxygen-deficient stratified reservoirs, hypolimnetic aeration or oxygenation and selective withdrawal can be evaluated, but their performance depends on reservoir geometry, operating conditions and sediment oxygen demand [4,10].

© 2026 London International Monthly Conference on Multidisciplinary Research and Innovation. All Rights Reserved.

Hypolimnetic oxygenation can help manage redox-sensitive metals, although sensor evidence shows that Fe and Mn concentrations may fluctuate at hourly-to-daily scales [3]. Managers should therefore monitor unintended effects, including mixing changes, energy consumption, downstream water temperature and water-quality trade-offs. Decision support should compare multiple scenarios rather than optimizing a single indicator.

4.2. Integrated adaptive management framework

An operational framework can proceed through five linked steps: establish a baseline and seasonal profile; identify dominant pressure sources; integrate satellite and sensor information; test interventions with calibrated models; and evaluate outcomes against measurable, reservoir-specific targets. A tiered monitoring network should cover inflow zones, the deepest basin and withdrawal structures. Sampling frequency should increase during heat waves, high-flow events, algal bloom periods and autumn turnover.

Table 3. Risk-based management actions and verification metrics.

Observed risk	Candidate intervention	Verification metric
High nutrient loading	Catchment nutrient control	TN, TP, Chl-a trends
Deep-water hypoxia	Oxygenation / withdrawal review	DO profiles and hypoxic duration
High suspended sediment	Erosion and inflow controls	TSS, turbidity, transparency
Fe/Mn mobilization	Redox and oxygen management	Depth-specific Fe, Mn, DO
Uncertain forecasts	Hybrid model + validation	Independent error metrics

The reviewed literature also reveals a persistent integration gap between water-quantity and water-quality observations [11]. Water-level and inflow data should therefore be analysed jointly with water chemistry and biological response. Machine-learning and Internet-of-Things systems can support automated warnings, but sensor reliability, explainability and data governance remain practical concerns [12].

4.3. Research gaps and implications

Three research gaps are especially important. First, optical remote sensing remains more mature for surface chlorophyll-a and suspended particles than for directly inferring dissolved nutrients or deep-water oxygen. Second, algorithms calibrated in one reservoir may fail under different optical, hydrological or climatic conditions. Third, management assessments often lack comparable cost, ecological side-effect and uncertainty metrics. These gaps support combined observational and process-based approaches rather than a single universal algorithm [5–9,11,12].

For semi-arid and strongly seasonal basins, including many Central Asian settings, a practical research design would pair monthly year-round profiles with more frequent warm-season sampling, event-based inflow sampling, and cloud-screened Sentinel-2/Landsat observations. These are proposed design choices, not measurements from a specific Central Asian reservoir. Thresholds for intervention should be established from applicable national standards, reservoir use and local baseline conditions.

5. Conclusion

Reservoir water quality is dynamic in three dimensions and through time. Recent evidence emphasizes the importance of stratification, oxygen depletion, sediment interactions and hydrological operations in shaping ecological and supply-water risks. Satellite remote sensing efficiently extends spatial coverage for optical indicators, while in situ depth profiles and laboratory analyses remain indispensable for nutrients, dissolved oxygen and metals. Machine-learning methods offer strong predictive potential within observed conditions, but physically based models are valuable for testing

© 2026 London International Monthly Conference on Multidisciplinary Research and Innovation. All Rights Reserved.

management interventions and unfamiliar future climates. A robust management strategy should combine upstream pollution prevention, risk-based vertical monitoring, integrated quantity–quality data and iterative evaluation of interventions. Future work should prioritize independent validation, uncertainty analysis and locally calibrated decision thresholds.

Declarations

Study type: narrative literature review. Data availability: no original dataset was generated. Funding: not specified. Conflict of interest: none declared for this manuscript draft. Author details and journal formatting to be completed before submission.

References

[1] Burkholder, J. M., et al. (2022). Classic indicators and diel dissolved oxygen versus trend analysis in assessing eutrophication of potable-water reservoirs. Ecological Applications, 32, e2541. https://doi.org/10.1002/eap.2541

[2] Esmaeilzadeh Hanjaniamin, A., Sarai Tabrizi, M., & Babazadeh, H. (2023). Dissolved oxygen concentration and eutrophication evaluation in Yamchi dam reservoir, Ardabil, Iran. Applied Water Science, 13, 9. https://doi.org/10.1007/s13201-022-01786-1

[3] Hammond, N. W., et al. (2023). High-frequency sensor data capture short-term variability in Fe and Mn concentrations due to hypolimnetic oxygenation and seasonal dynamics in a drinking water reservoir. Water Research, 240, 120084. https://doi.org/10.1016/j.watres.2023.120084

[4] Mi, C., Rinke, K., & Shatwell, T. (2024). Optimizing selective withdrawal strategies to mitigate hypoxia under water-level reduction in Germany's largest drinking water reservoir. Journal of Environmental Sciences. https://doi.org/10.1016/j.jes.2023.06.025

[5] Bangira, T., Matongera, T. N., Mabhaudhi, T., & Mutanga, O. (2024). Remote sensing-based water quality monitoring in African reservoirs, potential and limitations of sensors and algorithms: A systematic review. Physics and Chemistry of the Earth. https://doi.org/10.1016/j.pce.2023.103536

[6] Mi, C., Tilahun, A. B., Flörke, M., Dürr, H. H., & Rinke, K. (2024). Climate warming effects in stratified reservoirs: Thorough assessment for opportunities and limits of machine learning techniques versus process-based models in thermal structure projections. Journal of Cleaner Production. https://doi.org/10.1016/j.jclepro.2024.142347

[7] Mamun, M., Hasan, M., & An, K.-G. (2024). Advancing reservoirs water quality parameters estimation using Sentinel-2 and Landsat-8 satellite data with machine learning approaches. Ecological Informatics. https://doi.org/10.1016/j.ecoinf.2024.102608

[8] Sun, Y., Wang, D., Li, L., Ning, R., Yu, S., & Gao, N. (2024). Application of remote sensing technology in water quality monitoring: From traditional approaches to artificial intelligence. Water Research. https://doi.org/10.1016/j.watres.2024.122546

[9] Estimating the vertical profile of water quality variables in reservoirs: Application of remotely sensed data and machine learning techniques. (2025). Science of the Total Environment. https://doi.org/10.1016/j.scitotenv.2024.177543

[10] Hypolimnetic aeration optimization based on reservoir thermal stratification simulation. (2023). Journal of Hydrology, 625, 130106. https://doi.org/10.1016/j.jhydrol.2023.130106

[11] Ellis et al. (2024). Bridging the divide between inland water quantity and quality with satellite remote sensing: An interdisciplinary review. WIREs Water. https://doi.org/10.1002/wat2.1725

[12] A review of machine learning and internet-of-things on the water quality assessment: Methods, applications and future trends. (2025). Results in Engineering, 26, 105182. https://doi.org/10.1016/j.rineng.2025.105182

© 2026 London International Monthly Conference on Multidisciplinary Research and Innovation. All Rights Reserved.