PREVENTING GINGIVITIS IN ARTERIAL HYPERTENSION WHILE PRESERVING SALIVARY PROTECTIVE FUNCTIONS

Salomova Dilnora Abdimaxmudovna
Assistant, Department of Dentistry, Zarmed University
https://doi.org/10.5281/zenodo.23697773

Introduction

A preventive plan for gingivitis in a patient with arterial hypertension must be practical enough to sustain between visits. The plan also needs to account for dry-mouth complaints, changes in gingival contour, and the use of several oral care products. These factors can make cleaning uncomfortable or ineffective even when the patient reports brushing regularly. Recording what the patient actually does is therefore more informative than documenting advice alone.

Salivary protection includes the conditions that allow oral tissues and microbial communities to function. A preventive intervention should have a defined target, such as inaccessible plaque or persistent inflammation. Its value can then be assessed against that target and against tolerability. The objective of this synthesis is to organize these decisions into a reproducible preventive approach for adults with hypertension.

Materials and methods

A targeted narrative comparison considered gingivitis management guidance, the European consensus on plaque control, a randomized study of antibacterial mouthwash in treated hypertensive adults, and evidence on medication-associated gingival enlargement [1, 2, 3, 4]. The synthesis distinguished local outcomes from systemic observations. The published systolic blood pressure estimate and confidence interval were extracted from the mouthwash study for graphical presentation [2].

Clinical application was organized around four questions: which surfaces remain unclean, which local factors retain plaque, whether medication effects contribute to the gingival condition, and what specific benefit is expected from an adjunctive product. These questions informed a care matrix and a set of follow-up criteria. No pooled treatment effect was calculated.

Results

The proposed programme begins with observation of the patient's cleaning technique and identification of missed surfaces. Professional cleaning is directed to deposits and sites that prevent effective self-care. Interdental devices are selected according to access and comfort, with practical instruction [1, 3]. The next visit should establish whether the recommended technique is usable in the patient's daily routine, rather than merely whether the recommendation was remembered.

Medication review becomes relevant when enlargement of the gingiva changes access for cleaning. A systematic review identified calcium-channel blockers among medicines associated with this condition [4]. The dental record should describe onset, affected sites, and plaque retention. Discussion with the prescribing clinician may clarify the medication contribution, while local preventive care proceeds according to the observed tissue condition.

The mouthwash study provides a quantitative reason to consider the physiological consequences of adjunctive antisepsis. Three days of antibacterial mouthwash use increased systolic blood pressure by 2.3 mmHg relative to control, with a 95% confidence interval of 0.5–4.0 mmHg [2]. Figure 1 displays this

© 2026 London International Monthly Conference on Multidisciplinary Research and Innovation. All Rights Reserved.

estimate. The finding concerns a short exposure under study conditions and should not be interpreted as the effect of every mouthwash formulation or as a long-term cardiovascular outcome.




Figure 1. Published systolic blood pressure difference after three days of antibacterial mouthwash use in treated hypertensive adults [2]. The vertical error bar represents the 95% confidence interval, 0.5–4.0 mmHg. The column shows the estimated difference of 2.3 mmHg.

In the proposed approach, an adjunctive product is linked to a documented purpose and a planned review. The record specifies why mechanical care alone is temporarily insufficient, which outcome will indicate benefit, and which symptoms will prompt reconsideration. This makes a finite course distinguishable from indefinite self-directed use. The choice of formulation and duration follows the clinical indication and product instructions.

Table 1. Preventive actions and criteria for review

Care component	Intended purpose	Review criterion
Cleaning instruction	Reach missed surfaces	Technique and residual plaque
Professional care	Remove local deposits	Access and tissue response
Medication review	Clarify gingival enlargement	Onset and prescribing history
Adjunctive antisepsis	Address a defined short-term need	Benefit and tolerability
Salivary assessment	Investigate persistent dryness	Symptoms and secretion

Discussion

Preserving salivary protective functions requires attention to what prevention changes in the mouth. A fall in visible plaque is valuable, but an intervention that produces persistent discomfort may be abandoned. Follow-up should therefore include the patient's experience of brushing, interdental cleaning, taste, and dryness. These observations can identify a practical reason for an apparently poor clinical response.

© 2026 London International Monthly Conference on Multidisciplinary Research and Innovation. All Rights Reserved.

The distinction between a local endpoint and a systemic observation is central to interpretation. Gingival bleeding and plaque describe the periodontal response. Blood pressure measurements belong to cardiovascular assessment and cannot establish that a gingival intervention has succeeded or failed. If a relevant change is observed during use of an adjunctive product, its timing and circumstances should be communicated to the treating clinician without changing prescribed antihypertensive medication independently.

The proposed programme also separates a need for salivary investigation from a need for additional antimicrobial treatment. Persistent dryness may justify assessment of secretion and medication history. A biochemical deviation without an explanatory clinical pattern should first be checked for sampling and analytical variability. Treatment escalation is most defensible when the clinical problem, the intended action, and the expected outcome are explicitly connected.

The main limitation is that the reviewed evidence addresses different populations and interventions. A short mouthwash trial does not define the best long-term preventive regimen for all hypertensive patients. Future evaluation of this care approach should record baseline gingival status, medication exposure, adherence, and adverse experiences using consistent definitions. Such evaluation would establish whether structured review improves both inflammation control and the feasibility of daily care.

Conclusions

Prevention of gingivitis in arterial hypertension should combine accessible mechanical cleaning, correction of plaque-retentive factors, and review of medication-related oral changes. Adjunctive antisepsis is selected for a defined clinical purpose and reassessed for benefit and tolerability. Salivary investigations contribute when they address a specific unresolved question. This approach connects local disease control with preservation of a sustainable and physiologically informed care routine.

References

1. Chapple I.L.C., Van der Weijden F., Doerfer C. et al. Primary prevention of periodontitis: managing gingivitis. J Clin Periodontol. 2015;42(Suppl 16):S71–S76. doi:10.1111/jcpe.12366.
2. Bondonno C.P., Liu A.H., Croft K.D. et al. Antibacterial mouthwash blunts oral nitrate reduction and increases blood pressure in treated hypertensive men and women. Am J Hypertens. 2015;28(5):572–575. PMID:25359409.
3. Scottish Dental Clinical Effectiveness Programme. Gingivitis. Prevention and Treatment of Periodontal Diseases in Primary Care. https://periodontalcare.sdcep.org.uk/guidance/managing-disease/gingivitis/ (accessed 26 October 2026).
4. Nibali L., Lu E.M.C., Paz H.E.S. et al. Risk Factors and Management of Gingival Enlargement: A Systematic Review and Meta-Analysis. J Clin Periodontol. 2026. doi:10.1111/jcpe.70125.

© 2026 London International Monthly Conference on Multidisciplinary Research and Innovation. All Rights Reserved.