Email:

conference@houghtonstreetreview.com

Website:

www.houghtonstreetreview.com/index.php/conference

© 2026 International Online Conference on Multidisciplinary Academic Research. All Rights Reserved.

Editor-in-Chief:

Dr. Ewelum Johnson Nnadi

Managing Editors:

1.Harold Jan R. Terano

2.Dr. Mandakini Singla

3.Massoud Kaykhaii







COMPARATIVE PHYTOCHEMICAL AND BIOLOGICAL PERSPECTIVES ON PHLOMOIDES ISOCHILA AND PHLOMOIDES CANESCENS

Sotiboldiyev Akramjon Akbarali o‘g‘li

Abstract

Medicinal plants of the Lamiaceae family are valuable sources of structurally diverse natural compounds, but the chemical profiles and biological properties of individual species require species-specific verification. This thesis presents a comparative literature-based overview of Phlomoides isochila and Phlomoides canescens, focusing on reported phytochemical constituents and research directions for evaluating their biological potential. Published data on P. isochila describe an essential-oil profile in which β-thujone, camphor, and 1,8-cineole were reported as major volatile constituents. In vitro screening has also reported inhibition zones against selected Gram-positive bacteria, including Bacillus subtilis and Staphylococcus aureus. Research on P. canescens has highlighted polysaccharide fractions, including water-soluble carbohydrates and pectic or hemicellulose-related components, with attention to their interactions with microorganisms and possible prebiotic properties. These findings suggest that the two species may be investigated through complementary analytical strategies: gas chromatography–mass spectrometry for volatile compounds, liquid chromatography-based methods for non-volatile metabolites, and structural methods for polysaccharides. Biological evaluation should include standardized antioxidant and antimicrobial assays, appropriate positive and negative controls, concentration-dependent testing, and safety assessment. The available evidence does not justify assuming that compounds reported in other Phlomoides species occur in either target species without direct analytical confirmation. A coordinated comparative study using authenticated plant material and harmonized protocols would provide a stronger basis for characterizing their chemical diversity and assessing the relevance of their bioactive fractions for further research.

Keywords: Phlomoides isochila; Phlomoides canescens; Lamiaceae; phytochemical profile; essential oil; polysaccharides; antimicrobial screening; prebiotic potential.

Introduction

Natural products remain an important starting point for research in phytochemistry, pharmacognosy, and bioactivity screening. Members of the genus Phlomoides have been investigated for several classes of secondary metabolites, although chemical composition can vary among species and with ecological and methodological factors. Consequently, genus-level reports should guide research hypotheses rather than be treated as proof of a compound’s presence in a particular species.

Email:

conference@houghtonstreetreview.com

Website:

www.houghtonstreetreview.com/index.php/conference

© 2026 International Online Conference on Multidisciplinary Academic Research. All Rights Reserved.

Editor-in-Chief:

Dr. Ewelum Johnson Nnadi

Managing Editors:

1.Harold Jan R. Terano

2.Dr. Mandakini Singla

3.Massoud Kaykhaii







Aim

The aim is to compare the main phytochemical and biological research findings available for P. isochila and P. canescens and to outline an integrated methodology for their further investigation.

Materials and Methods

This thesis is based on a focused review of published literature and does not report newly performed experiments. For future comparative work, plant identity should be confirmed by a qualified botanist and voucher specimens deposited in a recognized collection. Collection site, season, developmental stage, drying procedure, and storage conditions should be documented. GC–MS is suitable for profiling volatile constituents; HPLC or LC–MS/MS can be used to investigate phenolic and other non-volatile metabolites; and carbohydrate analysis, molecular-mass determination, FTIR, or NMR may support characterization of polysaccharide fractions. Antioxidant activity may be screened using DPPH or ABTS assays, while antimicrobial testing should include standardized procedures and, where appropriate, MIC/MBC determination.

Key Findings from the Literature

For P. isochila, a published essential-oil study reported β-thujone, camphor, and 1,8-cineole among the principal volatile constituents. The same research reported in vitro inhibition zones against B. subtilis and S. aureus. Such measurements indicate activity under the specific test conditions and should not be interpreted as evidence of clinical effectiveness. The toxicological profile of individual constituents, particularly β-thujone, also needs to be considered before any application is proposed.

For P. canescens, published research has examined polysaccharide fractions and their interactions with microorganisms. Reported observations concerning beneficial microorganisms provide a rationale for further prebiotic-oriented investigations, but in vitro findings alone cannot establish benefits to human health. Detailed characterization of monosaccharide composition, molecular-mass distribution, and functional properties would help clarify which structural features may be associated with observed biological effects.

Discussion and Research Outlook

The available literature points to complementary research priorities for the two species: detailed essential-oil profiling and safety assessment for P. isochila, and structural characterization of polysaccharides and standardized prebiotic testing for P. canescens. A direct comparison should use the same sampling framework, extraction controls, analytical validation, and statistical procedures. Future work should also examine phenolic compounds and other metabolite groups in both species rather than extrapolating findings from related taxa. Replicated concentration-response experiments and appropriate reference standards are necessary to make results reproducible and comparable across laboratories.

Conclusion

Phlomoides isochila and Phlomoides canescens represent relevant subjects for comparative phytochemical research because the literature highlights different but potentially complementary fractions: volatile constituents in P. isochila and polysaccharides in P. canescens. Current findings provide a rationale for further

Email:

conference@houghtonstreetreview.com

Website:

www.houghtonstreetreview.com/index.php/conference

© 2026 International Online Conference on Multidisciplinary Academic Research. All Rights Reserved.

Editor-in-Chief:

Dr. Ewelum Johnson Nnadi

Managing Editors:

1.Harold Jan R. Terano

2.Dr. Mandakini Singla

3.Massoud Kaykhaii







investigation, not a basis for unqualified therapeutic claims. A systematic study combining validated chemical profiling, standardized bioassays, and safety evaluation is needed to determine the composition, reproducibility, and practical significance of the biological activities associated with these plants.

Selected References

1. Mohammadi, S., Jafari, B., Pourtaghi Anvarian, S., Nazemiyeh, H., Asnaashari, S., Delazar, A., & Asgharian, P. (2024). Taxonomic revision and clinical importance of Phlomoides genus: A comprehensive review. Pharmaceutical Sciences, 30(1), 21–35. https://doi.org/10.34172/PS.2023.1
2. Akramov, D. Kh., Jabbarov, Sh. A., Ashurova, L. N., & Akramov, T. R. (2023). Chemical composition and antimicrobial activity of the essential oil of the plant Phlomoides isochila. International Journal of Health Systems and Medical Sciences, 2(3), 73–76. https://doi.org/10.51699/ijhsms.v2i3.1307
3. Rahimova, H. R., & Alidjan, A. (2026). Phytochemical characteristics and biological activity of polysaccharide fractions of Phlomoides canescens. Universum: Chemistry and Biology, 6(144). https://7universum.com/en/nature/archive/item/22897