HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

FORENSIC MEDICAL THANATOLOGY: CONTEMPORARY PRINCIPLES, POSTMORTEM CHANGES, AND MEDICOLEGAL SIGNIFICANCE

X.V. Nurmatov
Assistant, Department of Pathological Anatomy and Forensic Medicine,

Andijan State Medical Institute; Independent Researcher,

Department of Forensic Medicine and Medical Law,

Tashkent State Medical University.

Abstract: Forensic medical thanatology is the branch of forensic medicine concerned with death, dying, postmortem transformation of the human body, determination of the cause and circumstances of death, and estimation of the postmortem interval. Its practical value lies in the fact that death initiates a sequence of physical, biochemical, cellular, and microbiological changes whose interpretation can provide important medicolegal information. The aim of this article was to synthesize the principal concepts of forensic thanatology and to evaluate the diagnostic value and limitations of early and late postmortem changes in contemporary forensic practice. A narrative integrative review was conducted using 15 national and international sources, including Uzbek forensic-medicine textbooks and major works in forensic pathology and postmortem science. The analysis showed that algor mortis, livor mortis, rigor mortis, ocular and drying changes, supravital reactions, autolysis, putrefaction, adipocere formation, mummification, and skeletal transformation must be interpreted as interdependent phenomena rather than isolated clocks. The reliability of postmortem interval estimation improves when scene information, body examination, temperature data, environmental conditions, autopsy findings, and laboratory methods are combined. Forensic thanatology also contributes to distinguishing natural from violent death, recognizing postmortem artifacts, evaluating body movement, selecting biological specimens, and reconstructing circumstances relevant to investigation. Modern imaging, biochemical assays, molecular techniques, and microbiome-based approaches broaden the field but do not eliminate the need for conventional morphological assessment. A standardized, multidisciplinary and cautious approach remains essential for scientifically defensible conclusions.

Keywords: forensic thanatology; postmortem interval; algor mortis; livor mortis; rigor mortis; decomposition; autolysis; forensic pathology; cause of death.

INTRODUCTION

Forensic medical thanatology is one of the fundamental sections of forensic medicine. It studies the biological and medicolegal aspects of death, the sequence of changes occurring in the body after

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

cessation of life, and the methods used to interpret these changes for legal and investigative purposes. Although death is an inevitable biological event, its forensic evaluation is complex because the human organism does not pass instantaneously from life to complete structural inactivity. Circulatory and respiratory arrest is followed by progressive loss of cellular function, redistribution of blood, biochemical exhaustion of energy reserves, changes in muscle tone, tissue dehydration, autolysis, microbial decomposition, and finally destruction or preservation of tissues depending on environmental conditions. Each stage may contain information useful to the forensic physician, but none should be interpreted outside the context in which the body was found.

The central medicolegal questions of thanatology include confirmation of death, determination of the cause and mechanism of death, estimation of the time elapsed since death, differentiation of injuries produced before death from changes occurring after death, assessment of whether a body has been moved, interpretation of postmortem artifacts, and selection of appropriate material for toxicological, histological, biochemical, microbiological, genetic, and other examinations. These questions are important in cases of violent death, sudden unexpected death, suspected poisoning, unidentified bodies, delayed discovery, exhumation, mass fatality events, and disputed medical circumstances. The quality of the answers depends on the integration of scene investigation, external examination, autopsy findings, laboratory studies, and knowledge of the person's medical history.

Traditional forensic thanatology has relied on visible and palpable signs such as cooling, hypostasis, rigor, drying, corneal changes, and decomposition. These signs remain indispensable because they can be assessed rapidly and at relatively low cost. However, their expression is strongly influenced by ambient temperature, clothing, body size, age, disease, blood loss, physical activity before death, humidity, airflow, immersion, burial, insect access, and many other variables. Therefore, modern forensic practice increasingly emphasizes a multimodal approach. Classical observations are combined with validated temperature models, chemistry of body fluids, histology, imaging, entomology, microbiology, and molecular analysis when appropriate.

The relevance of thanatology is especially high in healthcare systems because reliable determination and documentation of death support both justice and public health. An incorrect interpretation of postmortem change may lead to an erroneous conclusion about trauma, time of death, poisoning, neglect, or medical intervention. Conversely, a careful forensic examination can clarify whether an apparent lesion is a true injury or a postmortem artifact, whether lividity is compatible with the position in which the body was found, whether decomposition has obscured disease, and whether available evidence supports a narrow or broad estimate of the postmortem interval. The aim of this article was to summarize the contemporary principles of forensic medical thanatology, analyze the evidential value of major postmortem changes, and discuss their practical limitations and future directions.

MATERIALS AND METHODS

The study was designed as a narrative integrative review. Fifteen sources were selected to represent the principal areas of forensic thanatology: national forensic-medicine education in Uzbekistan, general forensic pathology, postmortem interval estimation, decomposition science, and environmental modification of postmortem processes. The national component included Uzbek-language forensic-medicine textbooks used in medical education. International sources included major

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

forensic pathology textbooks, monographs on human decomposition, and peer-reviewed publications addressing time since death and postmortem change.

The material was analyzed thematically rather than statistically. The review focused on six domains: the biological concept of death; early postmortem changes; supravital and biochemical phenomena; late postmortem transformation; estimation of the postmortem interval; and the medicolegal interpretation of thanatological findings. Priority was given to features that have direct practical relevance during scene examination, external examination, autopsy, exhumation, and laboratory assessment. Particular attention was paid to sources of error and to conditions that may accelerate, delay, imitate, or obscure postmortem processes.

Because the available literature includes different study designs and environmental settings, the analysis did not attempt to assign a universal numerical value to any single postmortem sign. Instead, findings were synthesized into a practical framework emphasizing combined interpretation. No experimental procedures were performed on human participants, no identifiable clinical data were used, and no new biological specimens were collected. The article therefore represents a literature-based scientific synthesis rather than an original autopsy or laboratory cohort study.

RESULTS

The review confirmed that forensic thanatology begins with a distinction between the event of death and the gradual postmortem extinction of cellular processes. Clinical observation may identify irreversible cessation of circulation and respiration, while specialized criteria are required in circumstances involving artificial support of vital functions. From a forensic perspective, the essential point is that after systemic death some tissues retain limited excitability for a period of time. This transient persistence explains supravital reactions and contributes to the development of rigor. It also means that the postmortem body is biologically dynamic: it continues to undergo predictable but highly variable physical and chemical changes after the loss of integrated organismal function.

Cooling of the body is one of the most familiar early postmortem phenomena. After heat production ceases, body temperature tends to approach environmental temperature through radiation, conduction, convection, and evaporation. The rate is not linear throughout the entire postmortem period. It is modified by ambient temperature, contact surface, air movement, clothing, blankets, body habitus, fever or hypothermia before death, immersion, and other circumstances. Rectal or deep body temperature can be useful when measured correctly and interpreted with a validated model, but a single temperature value cannot reliably define the time of death without scene information. Cooling is most useful when the environmental history is reasonably known and when measurements are obtained before substantial equilibration with the surroundings.

Livor mortis develops when circulation stops and blood settles under the influence of gravity in dependent vessels. It usually appears as patchy discoloration that becomes more extensive and eventually more fixed. Distribution of lividity may provide information about body position after death. Areas exposed to firm pressure may remain pale because vessels are compressed, creating contact pallor. If the body is moved before lividity becomes fixed, redistribution can occur; after fixation, the original pattern may persist and may coexist with a secondary pattern. This feature can support reconstruction of postmortem movement, although interpretation must consider blood loss, anemia, temperature, decomposition, pressure, and skin pigmentation. Color differences may raise

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

suspicion of particular toxic or environmental conditions, but color alone is insufficient for a definitive etiological diagnosis.

Rigor mortis reflects postmortem biochemical changes within skeletal muscle. Loss of adenosine triphosphate prevents dissociation of actin and myosin, resulting in progressive stiffening until proteolysis and decomposition later cause relaxation. The onset, intensity, distribution, and duration of rigor are influenced by temperature, muscle mass, premortem physical activity, convulsions, age, cachexia, and metabolic state. Because rigor can be broken mechanically and because decomposition can abolish it, its diagnostic value lies in the total pattern rather than in a rigid timetable. The coexistence of rigor with other early changes can help determine whether observations are consistent with the estimated interval, but isolated statements based solely on the presence or absence of rigidity are scientifically weak.

Postmortem drying and ocular changes form another group of early findings. Evaporation causes drying of exposed mucosa, the lips, areas of damaged epidermis, and the sclera when the eyelids remain partly open. Corneal clouding, loss of ocular tension, and other eye changes progress with time but depend strongly on exposure and humidity. Drying may darken abrasions and make them appear more prominent, creating a risk of confusing postmortem parchment-like change with true antemortem trauma. Recognition of these effects is important during external examination, especially in bodies discovered in dry, warm, or ventilated environments.

Supravital reactions represent residual tissue responsiveness after somatic death. Mechanical or electrical stimulation of muscle, pupillary responses to pharmacological agents, and localized contraction may persist for a limited period. Historically, such responses were used as auxiliary indicators of early postmortem interval. Their practical use is restricted by methodological variability and by the need for standardized stimulation and interpretation. Nevertheless, the underlying principle remains important: death of the organism does not imply simultaneous death of every cell. Biochemical changes in blood, vitreous humor, cerebrospinal fluid, muscle, and other tissues reflect this progressive loss of homeostasis and continue to be investigated as potential time-dependent markers.

Autolysis is driven primarily by endogenous enzymes and begins soon after cellular energy failure. It is particularly evident in organs rich in hydrolytic enzymes, including the pancreas and gastrointestinal tract, but it eventually affects all tissues. Autolytic changes can mimic disease and can complicate histological interpretation. Knowledge of expected postmortem morphology is therefore essential to avoid overdiagnosing inflammation, necrosis, or epithelial injury. Autolysis interacts with putrefaction, which is largely driven by microorganisms. After death, barriers that normally restrict microbial spread fail, allowing bacteria to proliferate and invade tissues. Gas production, discoloration, marbling of superficial vessels, skin slippage, bloating, purge fluid, and tissue liquefaction may follow in a sequence modified by the environment and the condition of the body.

Late postmortem change is particularly dependent on local conditions. Putrefaction generally progresses faster in warm and humid settings and slower under cold conditions. Access by insects and scavengers can markedly accelerate soft-tissue destruction. Burial limits some forms of access but introduces soil, moisture, depth, wrapping, and coffin characteristics as additional variables. Immersion changes heat loss, insect exposure, skin appearance, and microbial conditions. Extensive trauma may open new routes for microbial and insect activity. Obesity may promote heat retention and decomposition, whereas severe dehydration may favor preservation. Consequently, decomposition

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

stage can provide a broad indication of elapsed time but cannot be converted into a universal calendar interval without environmental reconstruction.

Two important preservative transformations are mummification and adipocere formation. Mummification results from rapid and substantial dehydration of tissues, producing dry, shrunken, leathery structures that may persist for long periods. It is favored by dry conditions, heat, low humidity, and air circulation, although combinations vary. Adipocere is a waxy alteration of body fat associated with hydrolysis and hydrogenation under moist conditions, often in water or damp burial environments. Both processes can preserve injuries, identity-related features, and other evidence that would otherwise be lost during ordinary putrefaction. Their presence is therefore not merely descriptive; it can substantially affect the strategy of examination and sampling.

Skeletonization represents an advanced endpoint of soft-tissue loss but may occur at highly variable rates. Once skeletal elements are exposed, forensic evaluation shifts toward the combined fields of forensic pathology, anthropology, odontology, and archaeology. Bones may retain evidence of perimortem trauma, disease, surgical intervention, age-related features, and identity. However, environmental damage, animal activity, root etching, weathering, burning, and postmortem breakage may imitate or obscure trauma. Thanatological interpretation remains relevant because the examiner must determine which changes belong to the period around death and which were created during the postmortem history of the remains.

The reviewed literature consistently supports a combined strategy for estimating the postmortem interval. In the early period, deep body temperature, lividity, rigor, ocular findings, and supravital reactions can be assessed together. As the interval increases, decomposition stage, insect development, environmental data, soil characteristics, body-fluid chemistry, microbial succession, and other methods become more important. Accuracy usually decreases as the interval widens because cumulative environmental variation becomes greater. The most defensible conclusion is often an interval or range accompanied by explicit conditions and limitations rather than a single precise time point.

Thanatology also plays a major role in determining cause, mechanism, and circumstances of death. The pattern of postmortem changes may help explain whether apparent hemorrhage is true blood extravasation or hypostasis, whether skin discoloration reflects injury or decomposition, whether fractures occurred around the time of death or later, and whether a body remained in one position. Autopsy adds internal evidence regarding disease, trauma, aspiration, blood loss, thrombosis, embolism, intoxication, and other causes. When decomposition is advanced, ancillary methods become increasingly important, but even extensively altered bodies may yield useful information through radiology, toxicology, histology, genetics, and careful anatomical dissection.

The overall result of the review is that no postmortem sign functions as an independent clock. Thanatological evidence is strongest when multiple findings are mutually consistent and when the examiner documents environmental conditions before interpreting them. A discrepancy between signs should not automatically be treated as error; it may indicate body movement, unusual premortem physiology, artificial heating or cooling, immersion, wrapping, medical intervention, or another modifying circumstance. The forensic value of postmortem change lies therefore not only in estimating time since death but also in revealing the postmortem history of the body.

DISCUSSION

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

The principal challenge in forensic medical thanatology is variability. Many classical descriptions were developed from repeated observations and remain clinically useful, but the biological processes themselves are continuous rather than divided into fixed time blocks. This distinction is important because legal questions often encourage requests for exactness that the science cannot always provide. A careful expert should distinguish direct observations from interpretation and should communicate uncertainty in a way that is understandable to investigators and courts. When environmental information is incomplete or contradictory, the width of the estimated postmortem interval should increase rather than being artificially narrowed.

Temperature-based estimation illustrates the importance of standardization. A useful measurement requires appropriate equipment, documentation of the measurement site and time, and a credible estimate of environmental conditions. Even then, factors such as clothing, body size, air movement, fever, hypothermia, and contact with heated or cooled surfaces can change heat transfer. Similar limitations apply to rigor and lividity. Their traditional time courses provide orientation, but they are not interchangeable with an objective clock. The best practice is to use these signs as components of a converging evidential framework.

Decomposition introduces even greater complexity because it results from interaction among endogenous autolysis, microorganisms, insects, scavengers, temperature, moisture, oxygen availability, burial conditions, and body-specific factors. Modern decomposition research has improved understanding of these processes and has encouraged quantitative methods such as accumulated thermal energy, entomological development models, chemical profiling, and microbial succession. These approaches are promising, but they require local validation. A model developed in one climate or burial environment may perform poorly in another. This is particularly relevant for countries with marked seasonal variation and diverse geographical conditions.

Laboratory approaches offer opportunities to complement gross examination. Biochemical investigation of vitreous humor and other fluids, postmortem changes in proteins and nucleic acids, metabolomic patterns, and microbial community shifts are active areas of research. Postmortem computed tomography and other imaging methods can document fractures, gas distribution, foreign bodies, and some internal pathology before dissection. These methods improve documentation and may reveal features that are difficult to appreciate macroscopically. However, advanced technology should be viewed as an extension of forensic reasoning rather than a replacement for it. Without accurate scene data and a complete morphological examination, highly technical results may still be misinterpreted.

A second major issue is the differentiation of true injury from postmortem artifact. Hypostasis can resemble bruising; skin slippage may resemble blistering; drying can emphasize abrasions; insect activity may create defects similar to wounds; postmortem fractures may be mistaken for perimortem trauma; and decomposition gas may distort tissues or produce unexpected radiological findings. Avoiding these errors requires knowledge of both injury biology and postmortem processes. When available, histology and targeted laboratory examination may assist, but the macroscopic distribution, anatomical relationships, hemorrhagic response, wound margins, and surrounding tissue changes remain central.

Forensic thanatology also has direct implications for specimen collection. The choice of samples depends on the stage of postmortem change and the investigative question. Peripheral blood may be preferable to central blood for some toxicological analyses because redistribution after death can alter

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

drug concentrations. Vitreous humor is relatively protected and may remain useful when blood is altered. Urine, bile, liver, muscle, hair, bone marrow, and other materials may become important in decomposed bodies. Samples for histology should include both lesions and appropriate comparison tissue. Genetic material can often be recovered long after soft-tissue decomposition. A thanatological assessment therefore guides not only interpretation but also the practical design of the laboratory investigation.

The educational importance of the field should be emphasized. Students and physicians need to understand that postmortem examination is not simply a list of signs to memorize. It is a structured reasoning process that begins at the scene and continues through external examination, autopsy, sampling, laboratory testing, synthesis, and reporting. Uzbek forensic-medicine literature provides a valuable national framework for teaching these fundamentals, while international research introduces evolving methods and broader comparative evidence. Combining both is important for training specialists who can work within national procedural requirements while applying contemporary scientific principles.

In practical forensic services, standardized documentation can substantially improve reliability. Body temperature, ambient temperature, clothing, coverings, position, degree of contact with surfaces, lividity distribution, rigor in different muscle groups, ocular state, drying, visible decomposition, insect activity, and other observations should be recorded systematically. Photographs and, where indicated, imaging provide a permanent record. The time at which each observation or specimen collection occurs should also be documented because postmortem changes continue during transport and examination. This is especially important when the initial scene is distant from the autopsy facility.

The narrative design of the present article is a limitation because it does not provide a pooled quantitative estimate of diagnostic accuracy for individual methods. In addition, environmental research from one region cannot be transferred directly to all other climates. Nevertheless, the consistency of the major principles across national textbooks, forensic pathology references, and decomposition research supports a clear practical conclusion: postmortem findings are most reliable when interpreted together, within their environmental and investigative context, and with explicit recognition of uncertainty.

CONCLUSION

Forensic medical thanatology provides the scientific framework for understanding death and the postmortem transformation of the human body. Early changes such as cooling, lividity, rigor, drying, ocular alterations, and supravital reactions are useful when assessed systematically, while autolysis, putrefaction, mummification, adipocere formation, and skeletonization become increasingly important as the postmortem interval lengthens. No single sign can determine time since death with universal precision. The most reliable medicolegal interpretation combines scene information, environmental conditions, external examination, autopsy, appropriate sampling, and laboratory methods. Modern imaging, biochemical, molecular, entomological, and microbiological techniques expand the possibilities of the discipline but should be integrated with classical morphological assessment. A standardized and cautious approach to thanatology strengthens cause-of-death investigation, reduces misinterpretation of postmortem artifacts, improves reconstruction of the postmortem history of the body, and supports scientifically defensible forensic conclusions.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

REFERENCES:

1. G‘iyosov, Z. A. (2018). Sud tibbiyoti [Forensic medicine]. Global Books.
2. Iskandarov, A. I., & Qo‘ldoshev, D. R. (2009). Sud tibbiyoti [Forensic medicine]. MERIYUS.
3. Indiaminov, S. I. (2012). Sud tibbiyoti [Forensic medicine]. Optima Print.
4. Saukko, P., & Knight, B. (2016). Knight’s forensic pathology (4th ed.). CRC Press.
5. Madea, B. (Ed.). (2014). Handbook of forensic medicine. Wiley-Blackwell.
6. Dolinak, D., Matshes, E. W., & Lew, E. O. (2005). Forensic pathology: Principles and practice. Elsevier Academic Press.
7. DiMaio, V. J. M., & DiMaio, D. (2001). Forensic pathology (2nd ed.). CRC Press.
8. Henssge, C., & Madea, B. (2007). Estimation of the time since death. Forensic Science International, 165(2–3), 182–184.
9. Megyesi, M. S., Nawrocki, S. P., & Haskell, N. H. (2005). Using accumulated degree-days to estimate the postmortem interval from decomposed human remains. Journal of Forensic Sciences, 50(3), 618–626.
10. Micozzi, M. S. (1991). Postmortem change in human and animal remains: A systematic approach. Charles C Thomas.
11. Goff, M. L. (2009). Early post-mortem changes and stages of decomposition in exposed cadavers. Experimental and Applied Acarology, 49, 21–36.
12. Campobasso, C. P., Di Vella, G., & Introna, F. (2001). Factors affecting decomposition and Diptera colonization. Forensic Science International, 120(1–2), 18–27.
13. Vass, A. A. (2001). Beyond the grave—understanding human decomposition. Microbiology Today, 28, 190–192.
14. Hayman, J., & Oxenham, M. (2016). Human body decomposition. Academic Press.
15. Prahlow, J. A. (2010). Forensic pathology for police, death investigators, attorneys, and forensic scientists. Humana Press.