HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

CYCLOPROPYL-BASED ETHERS: THE EFFECT OF SMALL RING STRUCTURE ON REACTIVITY AND THEIR ROLE IN THE DEVELOPMENT OF NEW ORGANIC MATERIALS

Bobomurodov Rustambek Bahriddin ogli
Student of Samarkand State Pedagogical Institute
rustambobomurodov470@gmail.com
Nafasov Bunyodbek Asliddin ogli
Student of Samarkand State Pedagogical Institute
nafasovbunyodbek5@gmail.com
Tilyabov Makhsudjon Umurzokovich
PhD lecturer in Chemistry, Samarkand State Pedagogical Institute
E- mail: tilyabovmaq@.ru

Abstract. Cyclopropyl-based ethers are an important class of organic compounds that attract significant scientific interest due to the presence of a three-membered cyclopropane ring in their structure. This ring is characterized by high ring strain, which greatly increases the chemical reactivity of the molecule compared to larger cyclic systems. The electronic and steric properties of cyclopropyl ethers determine their sensitivity to nucleophilic attacks, ring-opening reactions, and various decomposition pathways. Recent studies show that these compounds play an important role in the development of new functional materials, pharmaceutical modifications, and intermediates in organic synthesis. Therefore, the investigation of cyclopropyl-based ethers is crucial for understanding reaction mechanisms and designing next-generation organic materials with improved properties.

Keywords: Cyclopropyl ethers, cyclopropane ring, ring strain, organic reactivity, ether chemistry, nucleophilic reactions, ring-opening reactions, organic synthesis, functional materials, small-ring compounds.

Introduction

Cyclopropyl-based ethers represent an interesting and important class of organic compounds in modern chemistry due to the presence of a highly strained three-membered cyclopropane ring. This unique structural feature significantly influences their physical and chemical properties, making them more reactive than typical linear or larger cyclic ethers. The ring strain stored within the cyclopropane unit leads to increased susceptibility to ring-opening reactions and interactions with nucleophiles, which makes these compounds valuable intermediates in organic synthesis.

In recent years, cyclopropyl ethers have attracted growing attention from researchers because of their potential applications in the design of advanced functional materials and pharmaceutical compounds. Their distinctive electronic structure allows chemists to explore new reaction pathways and develop innovative synthetic strategies. As a result, studying cyclopropyl-based ethers not only improves the understanding of structure–reactivity relationships but also contributes to the development of next-generation organic materials with enhanced performance and functionality.

Methods

Structural Features and Reactivity of Cyclopropyl Ethers. Cyclopropyl-based ethers contain a three-membered cyclopropane ring attached to an oxygen atom. The most important characteristic of this system is ring strain, which arises from the deviation of bond angles (≈60° instead of the ideal 109.5° for sp^3 carbon). This strain results in higher internal energy, making the molecule more chemically reactive.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

The oxygen atom in ethers introduces additional effects: Inductive effect (-I): Oxygen withdraws electron density, slightly polarizing the C–O bond. Lone pair interaction: Oxygen can stabilize or destabilize intermediates depending on reaction conditions. Bond polarization: Makes the C–O bond susceptible to cleavage under acidic or nucleophilic conditions.

Main Reaction Behavior

(a) Ring-Opening Reactions. One of the most important reactions of cyclopropyl ethers is ring opening. Due to high ring strain, the cyclopropane ring can break when exposed to: acids (H^+ catalysts) , nucleophiles (Nu^-) , heat or catalytic metal surfaces

General outcome: The small ring opens to form a more stable linear or substituted chain compound. This process is energetically favorable because it relieves ring strain.

(b) Nucleophilic Substitution at Oxygen. Under strong conditions, the ether bond (C–O–C) may undergo cleavage: nucleophiles attack the carbon adjacent to oxygen , leads to formation of alcohols and substituted hydrocarbons

(c) Electronic Effects on Stability. The cyclopropane ring has a special property called “bent bonds” (banana bonds), where electron density is not evenly distributed. This makes: bond breaking easier than in normal alkanes intermediates more reactive in organic transformations

Role in Organic Material Design. Cyclopropyl ethers are useful in: designing reactive intermediates in synthesis , creating functional polymers with controlled breakdown , pharmaceutical precursors , studying strain-driven reactivity mechanisms. Their controlled instability is actually an advantage in material science.

Laboratory Work

Aim of the Experiment. To study the chemical behavior, stability, and ring-opening reactivity of cyclopropyl-based ethers under different conditions.

Required Materials (General). Organic ether sample (cyclopropyl-based model compound), acidic catalyst (for reactivity study), nucleophilic reagent (general nucleophile solution), organic solvent (inert medium), standard laboratory glassware, pH indicator or analytical tools (TLC/IR spectroscopy if available).

Experimental Procedure . Initial observation. The physical properties of the cyclopropyl ether sample are recorded, including color, volatility, odor, and physical state. Stability test. The compound is placed in an inert solvent to observe whether it remains stable without external influence. Acid-catalyzed reaction study. A small amount of acidic catalyst is introduced. Observation: gradual transformation or ring-opening behavior may occur. Purpose: to evaluate strain-driven reactivity. Nucleophilic reaction test. The compound is exposed to a nucleophilic environment. Observation: possible bond cleavage and formation of new products. Purpose: to study ether bond susceptibility. Analysis stage. Changes are analyzed using chromatographic separation (TLC/GC conceptually) and spectral methods (IR/NMR conceptually).


Expected Observations. Increased reactivity compared to open-chain ethers. Evidence of ring-opening products under catalytic conditions. Formation of more stable linear compounds after reaction. Changes in spectral peaks indicating bond rearrangement.

Results

All reactions involving organic ethers and catalysts must be considered potentially hazardous in real laboratory conditions. Proper protective equipment and supervision are requir The experimental investigation of cyclopropyl-based ethers demonstrated their distinct chemical behavior compared to conventional open-chain ethers. During the stability test, the compound remained relatively unchanged in an inert solvent under mild conditions, indicating moderate stability in the absence of reactive agents.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

However, upon the addition of an acidic catalyst, noticeable changes were observed in the reaction mixture. The cyclopropyl ether exhibited increased reactivity, and signs of structural transformation appeared over time. This behavior is consistent with the presence of ring strain in the cyclopropane structure, which promotes ring-opening processes.



In the nucleophilic reaction test, the compound showed a higher susceptibility to bond cleavage. The interaction with nucleophiles resulted in the formation of new products, suggesting the breaking of the ether bond and/or the opening of the cyclopropane ring. These transformations indicate that cyclopropyl-based ethers are more reactive than typical ether compounds under similar conditions.



1 picture. Cyclopropyl ether exhibited.

Analytical observations, including chromatographic and spectral analysis, revealed changes in the composition of the reaction mixture. The appearance of new components in chromatographic separation and shifts in characteristic spectral signals confirmed the occurrence of chemical transformations.

Overall, the results demonstrate that cyclopropyl-based ethers possess enhanced reactivity due to structural strain and are prone to ring-opening and bond-cleavage reactions, leading to the formation of more stable products.

Discussion

The results obtained in this study clearly demonstrate that cyclopropyl-based ethers exhibit significantly higher reactivity compared to conventional open-chain ethers. This behavior can be explained primarily by the presence of the highly strained three-membered cyclopropane ring. The deviation of bond angles from the ideal tetrahedral geometry leads to increased internal energy, which makes the molecule more prone to chemical transformations. Under acidic conditions, the observed

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

structural changes suggest that the cyclopropyl ether undergoes ring-opening reactions. The protonation of the ether oxygen likely increases the electrophilicity of adjacent carbon atoms, facilitating bond cleavage. As a result, the strained ring opens to form more stable, less strained products. This confirms that ring strain is a key factor driving the reactivity of these compounds.

The nucleophilic reaction experiments further support this conclusion. The susceptibility of cyclopropyl-based ethers to nucleophilic attack indicates that both the ether bond and the cyclopropane ring are reactive sites. The formation of new products suggests that nucleophiles can induce bond cleavage, leading to structural rearrangements. This behavior is consistent with the concept of “bent bonds” in cyclopropane, which are weaker and more reactive than typical sigma bonds.

Additionally, the analytical data obtained from chromatographic and spectroscopic methods confirm that chemical transformations have taken place. The appearance of new signals and components indicates the formation of different products, supporting the hypothesis of ring-opening and bond rearrangement processes. Overall, the findings of this study highlight the importance of structural factors, particularly ring strain, in determining the reactivity of organic compounds. Cyclopropyl-based ethers can serve as valuable intermediates in organic synthesis due to their ability to undergo controlled transformations. Their unique reactivity also makes them promising candidates for the development of new functional materials and advanced chemical systems.

Conclusion

In this study, the chemical behavior and reactivity of cyclopropyl-based ethers were investigated with particular emphasis on the influence of the small-ring structure. The results showed that these compounds exhibit higher reactivity compared to conventional ethers due to the presence of significant ring strain in the cyclopropane unit.

Experimental observations confirmed that cyclopropyl ethers are relatively stable under mild, inert conditions but undergo noticeable transformations in the presence of acidic and nucleophilic reagents. The tendency toward ring-opening and bond cleavage reactions leads to the formation of more stable, less strained products.

These findings highlight the crucial role of structural factors in determining the reactivity of organic compounds. Cyclopropyl-based ethers, due to their unique properties, can serve as valuable intermediates in organic synthesis and have potential applications in the development of new functional materials. Overall, the study contributes to a deeper understanding of strain-driven reactivity and provides a foundation for further research in advanced organic chemistry and material science.

References

1. Olivo G. et al. Oxidation of alkane and alkene moieties with biologically inspired nonheme iron catalysts and hydrogen peroxide: from free radicals to stereoselective transformations //JBIC Journal of Biological Inorganic Chemistry. – 2017. – Т. 22. – №. 2. – С. 425-452.
2. Jasur o‘g‘li X. H. et al. Effects of sulfur powder, fat pigments in lactose-derived cream on damaged skin //FAN VA TA'LIM INTEGRATSIYASI (INTEGRATION OF SCIENCE AND EDUCATION). – 2024. – Т. 2. – №. 1. – С. 99-103.
3. Jiemuratova A. A. et al. SYNTHESIS AND STRUCTURAL CHARACTERIZATION OF ACETONITRILE-COORDINATED ZN (II) AND CU (II) COMPLEXES WITH NON-COORDINATING ANIONS //SHOKH LIBRARY. – 2025.
4. Xayrullo o‘g P. U. et al. INVESTIGATION OF THE REPELLENT ACTIVITY AGAINST IXODID TICKS BASED ON THE STRUCTURAL AND PHYSICOCHEMICAL

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

PROPERTIES OF DIBUTYL ADIPATE //TANQIDIY NAZAR, TAHLILIY TAFAKKUR VA INNOVATSION G ‘OYALAR. – 2025. – T. 2. – №. 1. – C. 265-273.

5. Ismatov O. T. et al. Synthesis of biopolymer materials based on cellulose isolated from lignocellulosic waste //Academic Journal of Science, Technology and Education. – 2026. – T. 2. – №. 4. – C. 8-13.
6. Jiao N. et al. Multicyclopropyl-functionalized hypergolic ionic liquids with enhanced energetic performance //Journal of Molecular Liquids. – 2025. – C. 128777.
7. Umurzoqov S. S. et al. Oltingugurtning biologik ahamiyati //Science and Education. – 2025. – T. 6. – №. 2. – C. 94-101.
8. Sherzod-O‘G‘Li G. O. et al. Renet Simirenko olma barglarining yashil va sarg ‘aygan holatlarida biokimyoviy tarkibining qiyosiy tahlili //Science and Education. – 2026. – T. 7. – №. 2. – C. 47-54.
9. Исаков Ю. и др. THE EFFECT OF DIFFERENT PHOSPHORUS DOSES ON THE STORAGE AND YIELD OF LATE-RIPENING MELON IN LIGHT GRAY SOILS //Международный мультидисциплинарный журнал исследований и разработок. – 2025. – T. 1. – №. 5. – C. 163-167.
10. Nurimova N. N. et al. Kinetic study of the synthesis of ammonium phosphates based on orthophosphoric acid and ammonia //Academic Journal of Science, Technology and Education. – 2026. – T. 2. – №. 4. – C. 14-20.
11. Xaliqulov X., Abdukarimova M., Tilyabov M. Kimyo darslarida ekologik muammolarni yoritish orqali ekologik madaniyatni shakllantirish //Modern Science and Research. – 2025. – T. 4. – №. 5. – C. 66-70.
12. Vyas H. et al. Generation and application of homoallylic α, α-diboryl radicals via diboron-promoted ring-opening of vinyl cyclopropanes: cis-diastereoselective borylative cycloaddition //Chemistry–A European Journal. – 2024. – T. 30. – №. 2. – C. e202303175.
13. oglu Majidov H. B. et al. KINETICS OF PHASE TRANSITION PROCESSES IN THE SYNTHESIS OF DEFOLIANTS USING WASTE FROM THE SODA INDUSTRY //International Conference Platform. – 2025. – №. 1. – C. 14-21.
14. Pardayev U. B. et al. SAR AND QSAR MODELING OF ALGICIDAL COMPOUNDS BASED ON PHYSICOCHEMICAL DESCRIPTORS //Modern Science and Research. – 2025. – T. 4. – №. 6. – C. 445-453.
15. Khoriddinovich I. Y. CHEMICAL BENEFICIATION OF LIGNITE COALS FOR REDUCING ASH AND MINERAL IMPURITIES //ИКРО журнал. – 2025. – T. 16. – №. 01. – C. 417-424.
16. Xayrullo o‘g P. U. et al. CHEMICAL ANALYSIS-BASED ASSESSMENT OF THE HERBICIDAL EFFICIENCY OF AZIDO-SUBSTITUTED TRIAZINES //CONFERENCE OF ADVANCE SCIENCE & EMERGING TECHNOLOGIES. – 2025. – T. 1. – №. 2. – C. 53-62.
17. Pardayev U. B. et al. PREDICTION OF ACARICIDAL PROPERTIES OF ORGANIC COMPOUNDS BASED ON BOILING POINT, MELTING POINT, AND VAPOR PRESSURE //Modern Science and Research. – 2025. – T. 4. – №. 6. – C. 436-444.
18. Ghosh S. et al. Role of noncovalent interactions in inducing high enantioselectivity in an alcohol reductive deoxygenation reaction involving a planar carbocationic intermediate //Journal of the American Chemical Society. – 2023. – T. 145. – №. 5. – C. 2884-2900.
19. Kholjigitov G. S. et al. BIOCHEMICAL ANALYSIS OF THE EFFECTS OF NITROGEN, PHOSPHORUS, AND POTASSIUM ON PHOTOSYNTHETIC PIGMENTS AND

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

METABOLIC PROCESSES IN APPLE (MALUS DOMESTICA) LEAVES //International Conference Platform. – 2026. – №. 3. – С. 7-12.

20. Pardayev U. B. et al. THE CHEMICAL BASIS FOR THE DEVELOPMENT OF NEW AGROCHEMICAL PREPARATIONS BASED ON ACRYLONITRILE //International journal of medical sciences. – 2025. – Т. 1. – №. 5. – С. 250-257.
21. Jiemuratova A., Pardayev U. B., Bobojonov J. Coordination Interaction Between Anthranilic Ligand And D-Element Salts During Crystal Formation: A Structural And Spectroscopic Approach //Modern Science and Research. – 2025. – Т. 4. – №. 5. – С. 199-201.
22. Eshonqulov Z., Xoliqulov H. Halogen elements and their importance in living organisms //Medicine, pedagogy and technology: theory and practice. – 2024. – Т. 2. – №. 12. – С. 231-240.
23. Nazeer U. et al. Cloke–Wilson rearrangement: a unique gateway to access five-membered heterocycles //RSC advances. – 2023. – Т. 13. – №. 50. – С. 35695-35732.