HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

Neonatal Sepsis: Early Diagnosis, Risk Factors, and Modern Treatment Strategies

Davronova Samirabegim Umedovna
4th-year student, Faculty of General Medicine,
Samarkand State Medical University, Samarkand, Uzbekistan.
Rakhmatov Jasurbek Khurshid o'g'li
6th-year-student, Faculty of Pediatrics,
Samarkand State Medical University, Samarkand, Uzbekistan.
Mukhammadieva Barchinoy Mansur qizi
Rayimova Sabina Isroilovna
Matniyozova Parizoda Alijanovna
3rd-year students, Faculty of Pediatrics,
Samarkand State Medical University, Samarkand, Uzbekistan
doi:https://doi.org/10.5281/zenodo.24270739

Abstract: Neonatal sepsis is a serious systemic infection occurring during the first 28 days of life and remains one of the major causes of neonatal morbidity and mortality worldwide. It is particularly dangerous in premature and low-birth-weight infants because of their immature immune systems and limited physiological reserves. The disease may progress rapidly, leading to respiratory failure, septic shock, multiple organ dysfunction, and death. Early recognition and timely treatment are therefore essential for improving neonatal outcomes. The most important risk factors include prematurity, low birth weight, maternal infections, prolonged rupture of membranes, chorioamnionitis, invasive medical procedures, and prolonged hospitalization. Clinical manifestations are often nonspecific and may include temperature instability, respiratory distress, poor feeding, lethargy, apnea, and cardiovascular instability. Diagnosis is based on clinical assessment, blood culture, laboratory investigations, and evaluation of possible organ dysfunction. Modern treatment involves prompt empirical antibiotic therapy, appropriate supportive care, respiratory and hemodynamic stabilization, and subsequent adjustment of antimicrobial treatment according to microbiological results. This article reviews the major risk factors, clinical manifestations, diagnostic approaches, and contemporary treatment strategies for neonatal sepsis.

Keywords: neonatal sepsis, newborn, early-onset sepsis, late-onset sepsis, neonatal infection, blood culture, antibiotic therapy, premature infants, neonatal intensive care unit.

Introduction

Neonatal sepsis is a life-threatening condition caused by an invasive infection in newborn infants. It remains a significant challenge in neonatology because of its rapid progression, nonspecific clinical manifestations, and difficulties associated with early diagnosis.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

The neonatal period, defined as the first 28 days after birth, is characterized by significant physiological and immunological changes. Newborn infants, particularly those born prematurely, have immature immune responses and reduced ability to resist bacterial, viral, and fungal infections.

Neonatal sepsis is generally classified into early-onset sepsis and late-onset sepsis. Early-onset sepsis usually develops within the first 72 hours of life and is commonly associated with maternal infection and transmission of microorganisms before or during delivery. Late-onset sepsis occurs after 72 hours and is frequently associated with environmental exposure, healthcare procedures, and hospital-acquired infections.

The clinical course can vary from mild systemic manifestations to severe septic shock and multiple organ dysfunction. Because symptoms may initially be subtle, healthcare professionals must maintain a high level of clinical suspicion in newborns with recognized risk factors.

Advances in microbiological diagnostics, neonatal intensive care, antimicrobial treatment, and infection prevention have improved the management of neonatal sepsis. Nevertheless, early identification and appropriate treatment remain essential for reducing mortality and long-term complications.

Risk Factors and Etiology

The development of neonatal sepsis is influenced by maternal, perinatal, and neonatal factors. Prematurity and low birth weight are among the most important risk factors because immature immune function increases susceptibility to infection.

Maternal factors include group B streptococcal colonization, urinary tract infections, maternal fever during labor, chorioamnionitis, and prolonged rupture of membranes. These conditions increase the possibility of vertical transmission of microorganisms to the newborn.

Important neonatal risk factors include prematurity, very low birth weight, respiratory distress requiring mechanical ventilation, central venous catheterization, prolonged hospitalization, and surgical interventions.

The most common bacterial pathogens associated with early-onset sepsis include Group B Streptococcus and Escherichia coli. Other organisms may include Listeria monocytogenes and other Gram-negative bacteria.

Late-onset neonatal sepsis is frequently associated with coagulase-negative staphylococci, Staphylococcus aureus, Klebsiella pneumoniae, Escherichia coli, and other hospital-associated microorganisms. The distribution of pathogens varies according to geographical region, healthcare setting, and local antimicrobial resistance patterns.

Fungal infections, particularly those caused by Candida species, are important in extremely premature and critically ill newborns, especially those exposed to prolonged antibiotic treatment and invasive procedures.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

Clinical Manifestations

The clinical presentation of neonatal sepsis is often nonspecific, making early recognition difficult. Symptoms may develop gradually or progress rapidly over several hours.

Temperature instability is a common clinical feature. Infected newborns may develop fever or hypothermia, although normal body temperature does not exclude serious infection.

Respiratory manifestations include tachypnea, apnea, grunting, chest retractions, cyanosis, and increased oxygen requirements. Respiratory distress may be an early indication of systemic infection.

Neurological manifestations include lethargy, reduced activity, irritability, poor responsiveness, hypotonia, and seizures. Feeding difficulties, weak sucking, and refusal to feed are also important warning signs.

Gastrointestinal symptoms may include abdominal distension, vomiting, feeding intolerance, and diarrhea. Cardiovascular abnormalities include tachycardia, bradycardia, prolonged capillary refill time, hypotension, and poor peripheral perfusion.

Severe neonatal sepsis may progress to septic shock, disseminated intravascular coagulation, renal dysfunction, and multiple organ failure.

It is important to emphasize that no single clinical symptom is sufficiently specific to confirm or exclude neonatal sepsis. Clinical evaluation should always consider the infant's overall condition and the presence of maternal and neonatal risk factors.

Early Diagnosis and Laboratory Investigations

Early diagnosis is one of the most important factors influencing the prognosis of neonatal sepsis. Diagnostic evaluation should begin immediately when infection is suspected, without delaying necessary treatment in critically ill infants.

A detailed maternal and neonatal history is essential. Important information includes gestational age, birth weight, duration of membrane rupture, maternal temperature, maternal infections, delivery circumstances, and previous neonatal interventions.

Physical examination should assess respiratory function, heart rate, temperature, oxygen saturation, skin perfusion, neurological status, feeding ability, and urine output.

Blood culture remains the principal microbiological method for identifying the causative organism. An adequate blood sample should be collected before starting antibiotics whenever this can be done without delaying treatment.

A complete blood count may reveal leukopenia, leukocytosis, neutropenia, or an increased immature-to-total neutrophil ratio. However, these findings are not sufficiently sensitive or specific to establish the diagnosis independently.

C-reactive protein (CRP) is frequently used as an additional inflammatory marker. Serial measurements may help evaluate the inflammatory response and support decisions regarding the continuation of antibiotic therapy.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

Procalcitonin (PCT) is another biomarker that may assist in the assessment of neonatal bacterial infection. However, physiological changes during the first days of life must be considered when interpreting its results.

Lumbar puncture should be considered when meningitis is suspected, when blood cultures are positive, or when the infant's clinical condition indicates possible central nervous system infection. Cerebrospinal fluid examination may include cell count, glucose, protein, Gram staining, and culture.

Urine culture is particularly important in evaluating late-onset infection. Additional investigations, including blood gas analysis, glucose, electrolytes, renal and liver function tests, and coagulation studies, may be necessary depending on the severity of illness.

No laboratory marker should be interpreted in isolation. Diagnosis requires integration of clinical findings, microbiological results, and the infant's overall condition.

Modern Treatment Strategies

The treatment of neonatal sepsis requires rapid initiation of appropriate antimicrobial therapy and comprehensive supportive management. Newborns with suspected severe sepsis should be managed in a neonatal intensive care unit whenever possible.

Empirical antibiotic therapy

Empirical antibiotic treatment should be initiated promptly when neonatal sepsis is strongly suspected. The choice of antibiotics depends on the timing of infection, the likely pathogens, local antimicrobial resistance patterns, gestational age, and the infant's clinical condition.

For early-onset neonatal sepsis, a commonly recommended empirical combination is intravenous ampicillin and gentamicin. This combination provides coverage against important early-onset pathogens, including Group B Streptococcus, Listeria monocytogenes, and many susceptible Gram-negative organisms.

For late-onset sepsis, antimicrobial treatment should reflect local hospital microbiology and resistance patterns. Depending on the clinical setting, treatment may require coverage against staphylococci and Gram-negative bacteria.

When meningitis is suspected, the antimicrobial regimen must provide adequate penetration into the cerebrospinal fluid. Treatment should be selected according to the suspected pathogen and current neonatal protocols.

Once microbiological results become available, antibiotics should be adjusted according to the identified organism and its antimicrobial susceptibility. This approach reduces unnecessary broad-spectrum antibiotic exposure and helps prevent antimicrobial resistance.

Supportive and intensive care

Supportive treatment is essential in newborns with severe infection. Respiratory support may include supplemental oxygen, continuous positive airway pressure (CPAP), or mechanical ventilation, depending on the severity of respiratory failure.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

Hemodynamic stabilization involves careful assessment of circulation, perfusion, blood pressure, and urine output. Intravenous fluids may be administered when clinically indicated, with particular attention to avoiding fluid overload.

Vasoactive medications may be necessary in infants with persistent cardiovascular instability or septic shock.

Blood glucose, electrolyte balance, renal function, and acid-base status should be monitored regularly. Severe metabolic disturbances require appropriate correction.

Maintaining normal body temperature is also important because hypothermia and hyperthermia can worsen the metabolic demands of critically ill newborns.

Nutritional support should be individualized. Breast milk is preferred whenever clinically appropriate because it provides nutritional and immunological benefits. In unstable infants, feeding decisions should be based on gastrointestinal function and hemodynamic condition.

Prevention of Neonatal Sepsis

Prevention is a fundamental component of neonatal care. Effective infection-control practices can significantly reduce the risk of hospital-acquired infections.

Maternal screening and appropriate management of infections during pregnancy and labor are important preventive measures. Intrapartum antibiotic prophylaxis is recommended for eligible mothers at risk of transmitting Group B Streptococcus.

Strict hand hygiene, aseptic techniques, proper umbilical cord care, and careful management of intravenous catheters are essential in neonatal units.

Unnecessary invasive procedures and prolonged antibiotic exposure should be avoided. Breastfeeding and early skin-to-skin contact, when clinically appropriate, also contribute to improved neonatal health.

Healthcare professionals should follow standardized infection-prevention protocols and continuously monitor infection rates within neonatal intensive care units.

Prognosis and Long-Term Outcomes

The prognosis of neonatal sepsis depends on several factors, including gestational age, birth weight, causative microorganism, severity of infection, presence of organ dysfunction, and the time between symptom onset and treatment initiation.

Early recognition and appropriate antimicrobial therapy significantly improve the chances of survival.

However, severe neonatal sepsis may result in complications such as meningitis, neurological impairment, hearing loss, chronic lung disease, and developmental delay.

Premature infants and newborns with very low birth weight are particularly vulnerable to long-term complications. Regular developmental assessment and appropriate follow-up are therefore important for survivors of severe neonatal infections.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

Discussion

Neonatal sepsis remains a major clinical challenge because its initial symptoms are frequently subtle and nonspecific. The condition requires a combination of clinical experience, laboratory evaluation, microbiological investigation, and rapid therapeutic intervention.

Blood culture remains the principal diagnostic method, while inflammatory biomarkers such as CRP and procalcitonin provide additional information. Nevertheless, laboratory results should not delay treatment in infants with strong clinical suspicion of sepsis.

Modern neonatal management emphasizes appropriate empirical antibiotic selection, early respiratory and hemodynamic support, and regular reassessment of antimicrobial therapy. Antibiotics should be narrowed or discontinued when microbiological findings and the clinical course indicate that continued treatment is unnecessary.

Antimicrobial stewardship is especially important in neonatal intensive care units because excessive antibiotic exposure may contribute to antimicrobial resistance and disturbances in the developing intestinal microbiome.

Improving maternal care, strengthening infection-prevention practices, and ensuring access to neonatal intensive care are essential strategies for reducing the global burden of neonatal sepsis.

Conclusion

Neonatal sepsis is a potentially fatal infectious condition that requires early recognition, accurate diagnostic evaluation, and timely treatment. Prematurity, low birth weight, maternal infections, prolonged rupture of membranes, and invasive medical procedures are among the most important risk factors.

Clinical manifestations may include temperature instability, respiratory distress, poor feeding, lethargy, apnea, and cardiovascular abnormalities. Blood culture remains the primary microbiological diagnostic method, supported by laboratory investigations and careful clinical assessment.

Modern treatment combines prompt empirical antibiotic therapy, pathogen-directed antimicrobial treatment, respiratory support, hemodynamic stabilization, and appropriate nutritional management.

Prevention through maternal infection control, hand hygiene, aseptic procedures, and responsible antibiotic use is equally important.

Early diagnosis, evidence-based treatment, and multidisciplinary neonatal care remain the most effective approaches to reducing neonatal sepsis-related morbidity and mortality.

References

Uzbek Sources

1. Ministry of Health of the Republic of Uzbekistan. (2024). National Clinical Protocol for the Diagnosis and Treatment of Neonatal Sepsis. Tashkent, Uzbekistan.
2. Shamsiyev, A. M., & Mavlyanov, F. Sh. (2010). Neonatologiya [Neonatology]. Tashkent, Uzbekistan.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

3. Ismailov, S. I., & Rakhimov, A. R. (2018). Bolalar kasalliklari propedevtikasi [Propaedeutics of Childhood Diseases]. Tashkent, Uzbekistan.

4. International Sources

5. Stocker, M., et al. (2024). Management of neonates at risk of early onset sepsis: A probability-based approach and recent literature appraisal. European Journal of Pediatrics, 183, 5517–5529.
6. Shane, A. L., Sánchez, P. J., & Stoll, B. J. (2017). Neonatal sepsis. The Lancet, 390(10104), 1770–1780.