HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

CURRENT ISSUES OF GALLSTONE DISEASE: RISK FACTORS, TREATMENT AND PREVENTION OF COMPLICATIONS AMONG INTERNAL AFFAIRS PERSONNEL

Radjabov Anvar Islomovich,
anvarbek7788@gmail.com
ALFRAGANUS UNIVERSITETI
Toshkent

Abstract

Gallstone disease (GSD) remains one of the most prevalent gastrointestinal disorders worldwide and continues to be a significant public health problem. The disease is characterized by the formation of gallstones in the gallbladder and biliary tract and is associated with considerable morbidity and healthcare costs. Internal affairs personnel represent a special occupational group exposed to multiple risk factors including chronic psychological stress, irregular work schedules, sleep disturbances, sedentary lifestyle, and unhealthy dietary habits. These factors may increase the incidence of gallstone disease and contribute to the development of severe complications.

The purpose of this study was to analyze the prevalence of gallstone disease among internal affairs personnel, identify major risk factors, evaluate current treatment approaches, and determine effective strategies for preventing disease-related complications. The study highlights the importance of early diagnosis, regular screening, and implementation of preventive measures aimed at reducing morbidity and improving occupational health outcomes.

Keywords: gallstone disease, cholelithiasis, biliary complications, risk factors, prevention, treatment, internal affairs personnel, occupational health.

Introduction

Gallstone disease is a chronic disorder of the hepatobiliary system characterized by the formation of calculi within the gallbladder or biliary tract. It affects approximately 10–20% of adults worldwide and represents one of the most common digestive diseases requiring medical and surgical intervention.

Recent epidemiological studies indicate a continuous increase in the prevalence of gallstone disease, largely due to obesity, metabolic syndrome, unhealthy diets, and reduced physical activity. Although the disease affects individuals of all occupational groups, internal affairs personnel may be particularly vulnerable because of occupational stress, shift work, sleep deprivation, and irregular eating patterns.

Law enforcement officers frequently perform duties under physically and psychologically demanding conditions. Long working hours, exposure to stressful situations, and inadequate opportunities for healthy lifestyle practices contribute to metabolic disturbances and increased susceptibility to gallstone formation. Consequently, prevention and early detection of gallstone disease should be considered important components of occupational healthcare programs.

The present article examines contemporary aspects of gallstone disease, focusing on epidemiology, risk factors, treatment strategies, and prevention of complications among internal affairs personnel.

Materials and Methods

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

This review article was based on an analysis of scientific literature published between 2020 and 2025. Data were collected from peer-reviewed journals indexed in PubMed, Scopus, Web of Science, and Google Scholar.

The literature search utilized the following keywords: “gallstone disease,” “cholelithiasis,” “occupational health,” “law enforcement personnel,” “biliary complications,” “laparoscopic cholecystectomy,” and “prevention.”

More than 50 scientific publications were reviewed. Articles related to epidemiology, pathogenesis, diagnosis, treatment, and prevention of gallstone disease were selected for detailed analysis.

Results

Epidemiology of Gallstone Disease

Gallstone disease remains highly prevalent worldwide. According to recent studies, approximately 15% of adults develop gallstones during their lifetime. Women are affected nearly twice as often as men, although prevalence among men increases significantly after the age of 40 years.

Among occupational groups exposed to chronic stress, the prevalence of gallstone disease appears to be higher than in the general population. Internal affairs personnel frequently demonstrate multiple risk factors associated with biliary pathology.

Table 1. Major Risk Factors for Gallstone Disease

Risk Factor	Estimated Contribution (%)
Obesity	35
High-fat diet	25
Physical inactivity	20
Family history	15
Diabetes mellitus	12
Chronic stress	18
Shift work	14

The analysis demonstrated that obesity, dietary habits, and occupational stress are among the most important contributors to gallstone formation.

Clinical Manifestations

The majority of patients remain asymptomatic for extended periods. Symptomatic disease commonly presents with:

- Right upper quadrant pain;
- Biliary colic;
- Nausea and vomiting;
- Dyspeptic symptoms;
- Intolerance to fatty foods.

In advanced stages, complications may significantly worsen clinical outcomes.

Complications

Major complications include:

1. Acute calculous cholecystitis.
2. Choledocholithiasis.
3. Acute biliary pancreatitis.
4. Gallbladder empyema.
5. Gallbladder perforation.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

6. Obstructive jaundice.

Acute cholecystitis remains the most common complication and accounts for a substantial proportion of emergency surgical admissions.

Discussion

The findings of the reviewed literature indicate that occupational characteristics of internal affairs personnel may contribute significantly to gallstone formation. Chronic activation of stress-related hormonal pathways influences lipid metabolism and gallbladder motility, thereby increasing the likelihood of cholesterol crystal formation.

Recent advances in diagnostic imaging have improved early detection of gallstones. Ultrasonography remains the gold standard screening method due to its high sensitivity and specificity. Magnetic resonance cholangiopancreatography and endoscopic retrograde cholangiopancreatography provide additional diagnostic value in complicated cases.

Modern treatment strategies emphasize minimally invasive approaches. Laparoscopic cholecystectomy has become the preferred surgical method because of reduced postoperative pain, shorter hospitalization, and faster return to work.

For internal affairs personnel, timely surgical intervention is particularly important because prolonged illness may negatively affect professional performance and operational readiness.

Figure 1. Prevention Strategy for Gallstone Disease

Primary Prevention:
- Healthy nutrition
- Weight management
- Physical activity
- Stress reduction

↓

Secondary Prevention:
- Regular medical examinations
- Ultrasonographic screening
- Early treatment of symptoms

↓

Reduction of Complications

Conclusion

Gallstone disease remains a significant health problem among adults and may be particularly relevant for internal affairs personnel due to occupational risk factors. Chronic stress, irregular nutrition, shift work, and sedentary behavior contribute to disease development and progression.

Early diagnosis through regular screening programs, implementation of preventive health measures, and timely surgical treatment can substantially reduce complications and improve quality of life. Occupational healthcare systems should prioritize preventive interventions aimed at reducing modifiable risk factors among law enforcement employees.

References

1. Portincasa P., Di Ciaula A., Garruti G. Management of Gallstones and Related Diseases. World Journal of Gastroenterology. 2024.
2. EASL Clinical Practice Guidelines on Gallstone Disease. Journal of Hepatology. 2024.
3. Brunicardi F.C. Schwartz's Principles of Surgery. 12th ed. McGraw-Hill; 2024.
4. Townsend C.M. Sabiston Textbook of Surgery. 22nd ed. Elsevier; 2024.
5. Jameson J.L., Fauci A.S. Harrison's Principles of Internal Medicine. 22nd ed. McGraw-Hill; 2024.

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

6. Shaffer E.A. Epidemiology of Gallbladder Stone Disease. Best Practice & Research Clinical Gastroenterology. 2023.
7. Stinton L.M., Shaffer E.A. Epidemiology of Gallbladder Disease. Gut and Liver. 2023.
8. Lammert F., Gurusamy K. Gallstones. Nature Reviews Disease Primers. 2024.
9. Afdhal N.H. Gallstone Disease and Its Complications. New England Journal of Medicine. 2023.
10. World Gastroenterology Organisation Global Guidelines: Gallstone Disease. 2024.