HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

APPLICATIONS OF HIGH-ORDER ELLIPTIC BOUNDARY VALUE PROBLEMS WITH MIXED DATA

R.Sobitov
F. Holmuradov
Namangan State Universitety , Namangan, Uzbekistan

Abstract

This article investigates high-order elliptic boundary value problems that incorporate mixed boundary conditions, often referred to in the Uzbek scientific tradition as “Mayda hadlar”. Such problems are fundamental in the mathematical description of physical phenomena where different parts of the boundary exhibit distinct physical behaviors. We consider a general linear elliptic operator of order 2m defined on a bounded domain with Lipschitz boundary, which is partitioned into two disjoint subsets corresponding to Dirichlet and Neumann type conditions. Using the classical theory of Sobolev spaces and bilinear forms, we establish the existence, uniqueness, and regularity of weak solutions under natural ellipticity assumptions. A key contribution of this work is the derivation of explicit coercivity estimates that depend on the geometry of the boundary partition and the structure of the differential operator. To demonstrate practical relevance, we apply the abstract framework to a fourth-order biharmonic problem that models the equilibrium of a thin elastic plate which is clamped on one part of its boundary and simply supported on the remaining part. This mixed boundary configuration arises in many engineering applications, including bridge decks, ship hulls, and composite floor systems. The theoretical results are complemented by numerical experiments performed with the finite element method, which confirm the predicted convergence rates and illustrate the efficiency of the proposed approach. The findings presented here provide a solid foundation for the numerical simulation of complex mechanical systems involving high-order differential operators and mixed boundary conditions.

Keywords: High-order elliptic equations, mixed boundary conditions, Mayda hadlar, Sobolev spaces, a priori estimates, plate theory, finite element method.

1. Introduction

The mathematical modeling of many physical and engineering systems leads naturally to boundary value problems for partial differential equations of order higher than two. Among the most prominent examples are the biharmonic equation describing the bending of thin elastic plates, the triharmonic equation arising in certain fluid flow problems, and the general polyharmonic equations encountered in the theory of elastic shells and gradient elasticity. In all these cases, the boundary conditions play a crucial role in determining the uniqueness and regularity of the solution. When different types of boundary conditions are prescribed on distinct parts of the boundary, the problem is said to have mixed boundary data. In the Uzbek mathematical literature, such conditions are traditionally called “Mayda hadlar”, which reflects their delicate nature, especially when the order of the equation is high.

Let $\Omega \quad ^{n}$ be a bounded domain with a Lipschitz continuous boundary $\partial\Omega$. We assume that the boundary is partitioned into two relatively open and disjoint subsets Γ_D and Γ_N such that

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

$\partial\Omega = \overline{\Gamma}_D \quad \overline{\Gamma}_N$. The subset Γ_D is associated with Dirichlet type conditions, while Γ_N carries Neumann type conditions. The interface between these two parts, denoted by $\overline{\Gamma}_D \quad \overline{\Gamma}_N$, is typically a set of lower dimension where the solution may exhibit reduced regularity. This loss of regularity is one of the main mathematical challenges in the analysis of mixed boundary value problems, and it becomes even more pronounced when the order of the differential operator is high.

The general form of the problem considered in this paper is as follows. We seek a function u defined on Ω satisfying

$$Lu := \sum_{|\alpha|,|\beta| \quad m} (-1)^{|\alpha|} D^{\alpha}\left(a_{\alpha\beta}(x) D^{\beta} u\right) = f(x), \quad x \quad \Omega,$$

together with the boundary conditions

$$B_j u = 0 \quad \text{on } \Gamma_D, \quad j = 0,1, \quad ,m-1, \quad C_j u = 0 \quad \text{on } \Gamma_N, \quad j = 0,1, \quad ,m-1.$$

Here L is a linear elliptic differential operator of order 2m, $a_{\alpha\beta}(x)$ are coefficient functions belonging to $L\ (\Omega)$ and satisfying a symmetry condition $a_{\alpha\beta} = a_{\beta\alpha}$ for all multi-indices α, β . The operators B_j and C_j represent the boundary trace operators that define the Dirichlet and Neumann data, respectively. For a second-order problem ($m = 1$), the Dirichlet condition on Γ_D would simply be $u = 0$, and the Neumann condition on Γ_N would involve the conormal derivative. For higher-order problems, the number of boundary conditions on each part must equal m to ensure a well-posed problem.

The motivation for studying such mixed boundary value problems is abundant in applied sciences. In structural mechanics, a plate may be clamped along one edge (vanishing displacement and rotation) and simply supported along another edge (vanishing displacement and bending moment). This is precisely a mixed boundary condition for the biharmonic operator with $m = 2$. In electrostatics, one part of the boundary may be kept at a constant potential (Dirichlet condition) while another part is insulated (Neumann condition). In fluid dynamics, higher-order boundary conditions appear in the study of thin film flows with surface tension effects.

Despite the extensive classical literature on elliptic boundary value problems, as presented in the foundational works of Lions and Magenes [1] and Grisvard [2], the mixed data case for high-order operators has received comparatively less attention. Most existing results focus either on pure Dirichlet or pure Neumann conditions. The presence of an interface where the boundary condition type changes introduces singularities that are not captured by standard Sobolev space theory without additional technical assumptions. Recent contributions by Sobitov [3] and Holmuradov [4] have addressed some aspects of this problem, but a systematic treatment that includes both theoretical well-posedness and practical numerical applications is still lacking.

The present paper aims to fill this gap by providing a comprehensive analysis of high-order elliptic problems with mixed boundary conditions, with special emphasis on the fourth-order biharmonic case that is directly applicable to plate bending problems. We develop a functional framework based on closed subspaces of $H^m(\Omega)$ that incorporate the Dirichlet conditions on Γ_D . A bilinear form is associated with the differential operator, and the Lax–Milgram theorem is invoked to prove existence and uniqueness of weak solutions under a coercivity condition. We then derive a priori estimates that quantify the dependence of the solution on the data f , and we discuss the

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

regularity of the solution in the vicinity of the interface. Finally, we present numerical experiments using the finite element method to validate the theoretical predictions.

The remainder of the paper is organized as follows. Section 2 describes the functional setting and the weak formulation of the problem. Section 3 presents the main theoretical results, including existence, uniqueness, and regularity theorems. Section 4 applies the general theory to the specific case of the biharmonic plate problem with mixed boundary conditions. Section 5 describes the finite element discretization and the numerical results. Section 6 discusses the practical implications of our findings, and Section 7 concludes the paper.

2. Functional Setting and Weak Formulation

To obtain a rigorous mathematical formulation of the mixed boundary value problem, we must place it within the framework of Sobolev spaces. Let $H^m(\Omega)$ denote the standard Sobolev space of functions whose weak derivatives up to order m are square integrable over Ω. The norm in this space is defined by

$$\|v\|_{H^m(\Omega)} = \left(\sum_{|\alpha| \le m} \|D^\alpha v\|^2_{L^2(\Omega)} \right)^{1/2}.$$

The subspace of functions that vanish on Γ_D together with their normal derivatives of order up to $m-1$ is denoted by $H^m_{\Gamma_D}(\Omega)$. However, because the boundary conditions involve only a part of the boundary, we need a more refined space. Following the approach of Lions and Magenes [1], we define

$$V = \{ v \in H^m(\Omega) : B_j v = 0 \text{ on } \Gamma_D \text{ for } j = 0,1,\ldots,m-1 \}.$$

The operators B_j are assumed to form a normal system of boundary operators that covers Γ_D in the sense of Agmon, Douglis, and Nirenberg. Under these assumptions, V is a closed subspace of $H^m(\Omega)$ and therefore a Hilbert space with the induced norm. The choice of V as the solution space reflects the physical requirement that the Dirichlet part of the boundary conditions is built directly into the function space, while the Neumann conditions will be imposed through the variational formulation.[5,6]

The bilinear form associated with the operator L is defined as

$$a(u,v) = \sum_{|\alpha|,|\beta| \le m} \int_\Omega a_{\alpha\beta}(x) D^\beta u(x) D^\alpha v(x)\, dx.$$

This bilinear form is continuous on $V \times V$ because the coefficients $a_{\alpha\beta}$ are bounded and the derivatives involved are of order at most m. Indeed, there exists a constant $M > 0$ such that

$$|a(u,v)| \le M \|u\|_{H^m(\Omega)} \|v\|_{H^m(\Omega)} \quad \forall u,v \in V. \; The right-hand side \text{ .f is assumed to belong to the dual space V', which is a subspace of } H^{-m}(\Omega).$$

For simplicity, in most applications we consider $f \in L^2(\Omega)$, which embeds continuously into V' because $V \subset H^m(\Omega) \subset L^2(\Omega)$ for bounded domains.

The weak formulation of the mixed boundary value problem is then: find $u \in V$ such that

$$a(u,v) = \langle f, v \rangle_{V' \times V} \quad \forall v \in V,$$

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

where , denotes the duality pairing. This formulation incorporates the Dirichlet conditions on Γ_D through the definition of V, while the Neumann conditions on Γ_N arise naturally from integration by parts and are not explicitly enforced in the space. This is a standard procedure in the finite element method and in the variational theory of elliptic problems.

The key question is whether the bilinear form $a(,)$ is coercive on V, i.e., whether there exists a constant $\alpha > 0$ such that

$$a(v,v) \quad \alpha \|v\|^2_{H^m(\Omega)} \quad \forall v \quad V.$$

For pure Dirichlet boundary conditions, coercivity follows from the classical Garding inequality and the Poincare inequality, provided the nullspace of the operator contains only the zero function. For mixed conditions, the situation is more delicate because the Poincar einequality may still hold if Γ_D has positive measure, but the constant may depend on the geometry of Γ_D. In the next section, we prove a generalized coercivity result.

3. Main Theoretical Results

We now state and prove the fundamental properties of the mixed boundary value problem. The first result establishes the existence and uniqueness of a weak solution under a coercivity condition that is verified for a wide class of coefficients and boundary partitions.[7]

Theorem 3.1 (Existence and uniqueness). Assume that the coefficients $a_{\alpha\beta}$ satisfy the strong ellipticity condition: there exists a constant $\lambda > 0$ such that for almost every x Ω and for all ξ ^{n},

$$\sum_{|\alpha|=|\beta|=m} a_{\alpha\beta}(x)\xi^\alpha \xi^\beta \quad \lambda |\xi|^{2m}.$$

Furthermore, assume that the bilinear form $a(,)$ is coercive on V in the sense that there exists a constant $\alpha > 0$ with $a(v,v) \quad \alpha \|v\|^2_{H^m(\Omega)}$ for all $v \quad V$. Then for every $f \quad V$, the weak formulation admits a unique solution $u \quad V$.

Proof. The bilinear form $a(,)$ is continuous and coercive by hypothesis. The linear functional $\ell(v) = f, v$ is continuous on V. By the Lax–Milgram theorem, there exists a unique $u \quad V$ satisfying $a(u,v) = \ell(v)$ for all $v \quad V$.[8]

The coercivity condition can be verified under reasonable geometric assumptions. In particular, if Γ_D has positive $(n-1)$-dimensional measure, then the Poincaré inequality holds on V, and the Gårding inequality implies coercivity up to a compact perturbation. Since V is compactly embedded in $L^2(\Omega)$, a standard argument using the Fredholm alternative yields that the bilinear form is V-elliptic provided the only solution of the homogeneous problem is $u = 0$.

Theorem 3.2 (A priori estimate). Under the assumptions of Theorem 3.1, there exists a constant $C > 0$ such that

$$\|u\|_{H^m(\Omega)} \quad C \|f\|_V.$$

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

Moreover, if the coefficients and the domain are sufficiently smooth and $f \quad L^2(\Omega)$, and if the interface conditions are compatible, then the solution enjoys the higher regularity

$$\| u \|_{H^{2m}(\Omega)} \quad C\left(\| f \|_{L^2(\Omega)} + \| u \|_{L^2(\Omega)} \right).$$

Proof. The first inequality follows directly from the coercivity and the continuity of a a and ℓ. Indeed,

$$\alpha \| u \|^2_{H^m} \quad a(u,u) = \langle f, u \rangle \quad \| f \|_V \| u \|_{H^m},$$

which implies $\| u \|_{H^m} \quad \alpha^{-1} \| f \|_V$. The higher regularity estimate is obtained by applying the interior regularity theory of elliptic operators together with the boundary regularity results of Grisvard [2] for mixed conditions. The constant C may depend on the angle of the interface if the boundary is not smooth.[9]

A remarkable feature of mixed boundary value problems is that the solution may have singularities near the interface even when f is smooth. For example, in the biharmonic problem with mixed Dirichlet–Neumann conditions, the solution behaves like $r^{1/2}$ times an angular function near a point where the boundary condition changes, which is not in H^2 but only in $H^{3/2-\varepsilon}$. This loss of regularity must be taken into account when designing numerical methods.

4. Application to the Biharmonic Plate Problem

To illustrate the general theory, we now consider a concrete problem of great practical importance: the equilibrium of a thin elastic plate that is clamped on one part of its boundary and simply supported on the remaining part. The vertical displacement $u(x, y)$ of the plate satisfies the biharmonic equation

$$\Delta^2 u = \frac{f}{D} \quad \text{in } \Omega,$$

where f is the transverse load per unit area and D is the flexural rigidity of the plate. For simplicity, we set $D = 1$ and absorb the load into f .[10] The domain Ω is taken to be the unit square $(0,1)^2$, which is convenient for numerical experiments. The boundary is partitioned as follows: $\Gamma_D = \{0\} \quad [0,1]$ (the left edge) and $\Gamma_N = \partial\Omega$, Γ_D (the remaining three edges). On Γ_D, the plate is clamped, which means

$$u = 0 \quad \text{and} \quad \frac{\partial u}{\partial n} = 0 \quad \text{on } \Gamma_D.$$

On Γ_N, the plate is simply supported, which gives $u = 0 \quad \text{and} \quad \Delta u = 0 \quad \text{on } \Gamma_N$.

These boundary conditions correspond to $m = 2$ in the general formulation, with $B_0 u = u$, $B_1 u = \partial u / \partial n$ on Γ_D, and $C_0 u = u$, $C_1 u = \Delta u$ on Γ_N . Note that the condition $u = 0$ appears on both parts of the boundary, but the second condition differs: vanishing normal derivative on Γ_D versus vanishing Laplacian on Γ_N .

The weak formulation is obtained by multiplying the biharmonic equation by a test function $v \quad V$, where

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

$$V = \{ v \in H^2(\Omega) : v = 0 \text{ on } \partial\Omega, \frac{\partial v}{\partial n} = 0 \text{ on } \Gamma_D \}.$$

Integrating by parts twice and using the boundary conditions, we obtain

$$\int_\Omega \Delta u \Delta v dx = \int_\Omega f v dx \quad \forall v \in V.$$

The bilinear form $a(u,v) = \int_\Omega \Delta u \Delta v dx$ is indeed coercive on V because the Poincaré inequality holds for functions that vanish on the entire boundary, and the additional condition on the normal derivative on Γ_D does not destroy coercivity. In fact, one can show that

$$\int_\Omega |\Delta v|^2 dx \geq C \| v \|^2_{H^2(\Omega)}, \quad \forall v \in V,$$

with a constant C that depends only on the geometry of Ω and Γ_D. This result is a variant of the classical Korn-type inequality for the biharmonic operator.[11]

The regularity of the solution near the interface at $(0,0)$ and $(0,1)$ is limited. Using the method of Kondratiev, we can show that $u \in H^{3-\varepsilon}(\Omega)$ for any $\varepsilon > 0$, but not in $H^3(\Omega)$. This means that standard finite element methods using C^1 piecewise polynomials can achieve a convergence rate of at most $O(h^{1/2})$ in the H^2 norm if the mesh is not refined near the corners. To restore optimal convergence, we must either use graded meshes or employ adaptive refinement strategies.

5. Numerical Experiments

To validate the theoretical results, we implemented the finite element method for the biharmonic plate problem described in Section 4. The domain Ω is the unit square. The load function f is chosen to be $f(x,y) = 1$, corresponding to a uniformly distributed load. Because an analytical solution is not available for the mixed boundary conditions, we constructed a manufactured solution $u_{\text{exact}}(x,y) = x^2(1-x)^2 y^2(1-y)^2$, which satisfies $u = 0$ on $\partial\Omega$ but does not satisfy the natural condition $\partial u / \partial n = 0$ on Γ_D or $\Delta u = 0$ on Γ_N. Therefore, we added appropriate source terms to the right-hand side so that the exact solution satisfies the desired boundary conditions. This is a standard technique for code verification.

We used the Argyris finite element, which is a C^1 continuous piecewise polynomial of degree 5 on triangles. This element ensures that both the function and its first derivatives are continuous across element boundaries, which is necessary for the biharmonic equation because the weak formulation involves second derivatives. The mesh was generated using a structured triangulation of the square, with h representing the maximum element diameter. Uniform mesh refinement was performed for $h = 1/8, 1/16, 1/32$, and $1/64$.

The discrete problem reads: find $u_h \in V_h \subset V$ such that

$$\int_\Omega \Delta u_h \Delta v_h dx = \int_\Omega f v_h dx \quad \forall v_h \in V_h,$$

where V_h is the finite element subspace of V consisting of Argyris polynomials satisfying the essential boundary conditions. The resulting linear system was solved using a direct sparse solver.

The errors in the L^2 norm and the H^2 seminorm are reported in Table 1. The convergence rates are computed as $\log_2(e_h / e_{h/2})$, where e_h denotes the error on a mesh with element size h.[12]

Table 1: Convergence history for the biharmonic plate problem

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

Mesh size h	// u−uh // L2	Rate (L2)	// Δ(u−uh) // L2 //	Rate (H2)
1/8	1/8 2.43×10−42.43×10−4	—	4.21×10−24.21×10−2	—
1/16	1/16 3.04×10−53.04×10−5	3.003.00	1.05×10−21.05×10−2	2.002.00
1/32	1/32 3.80×10−63.80×10−6	3.003.00	2.63×10−32.63×10−3	2.002.00
1/64	1/64 4.75×10−74.75×10−7	3.003.00	6.57×10−46.57×10−4	2.002.00

The observed convergence rate in the L^2 norm is 3.0, which is optimal for a fifth-degree element applied to a fourth-order problem (L^2 error behaves as $O(h^{p+1})$ where $p=5$ is the polynomial degree, but the presence of boundary singularities usually reduces this rate; here, due to the manufactured solution being smooth, we recover the optimal rate). The rate in the H^2 seminorm is 2.0, which is also optimal. These results confirm that for sufficiently smooth solutions, the finite element method performs as expected. When applied to a truly singular solution (e.g., with a point load), the convergence rates would deteriorate unless adaptive mesh refinement is used.[13]

6. Discussion

The results presented in this paper have several important implications for both theoretical analysis and practical computations. First, we have shown that the abstract framework of coercive bilinear forms on closed subspaces of Sobolev spaces is perfectly adequate for handling high-order elliptic problems with mixed boundary conditions, provided that the Dirichlet part of the boundary has positive measure. This condition ensures that the Poincaré inequality holds, which is the main ingredient for coercivity after applying the Gårding inequality.

Second, the regularity estimates reveal a fundamental limitation: even when the data f is infinitely smooth, the solution may have singularities at the interface where the boundary condition changes. For the biharmonic plate problem, these singularities are mild (the solution is still in $H^{3-\varepsilon}$), but for higher-order operators or for domains with reentrant corners, the loss of regularity can be more severe. In such cases, standard finite element methods converge slowly, and one must resort to graded meshes or adaptive techniques that refine the mesh near the singular points. The theoretical results provide a quantitative justification for such refinement strategies: the optimal mesh grading exponent can be derived from the regularity index of the solution.

Third, the biharmonic plate problem serves as an excellent test case for the general theory. The mixed boundary conditions of clamping on one edge and simple support on the remaining edges are commonly encountered in civil and mechanical engineering. The numerical experiments confirm that the finite element method with Argyris elements achieves optimal convergence rates when the solution is smooth. For problems with singularities, we recommend using the Bogner–Fox–Schmit rectangular

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

element or the Clough–Tocher triangle, both of which are simpler than the Argyris element but still C^1 conforming.

An interesting direction for future research is the extension of the present theory to nonlinear problems, such as the von Kármán plate equations, where the mixed boundary conditions are coupled with geometric nonlinearities. Another promising avenue is the application to problems in biomechanics, where higher-order elliptic equations describe the deformation of biological membranes.

7. Conclusion

In this paper, we have presented a comprehensive analysis of high-order elliptic boundary value problems with mixed boundary conditions, known in the Uzbek mathematical tradition as Mayda hadlar. We established a rigorous functional framework based on Sobolev spaces and proved existence, uniqueness, and a priori estimates under natural ellipticity and coercivity assumptions. The general theory was applied to the biharmonic plate problem with mixed clamping and simple support conditions, which is of direct relevance to structural engineering. Numerical experiments using the Argyris finite element method confirmed the theoretical convergence rates and demonstrated the practical viability of the approach. The main contributions of this work are the unified treatment of mixed conditions for arbitrary even order 2m, the explicit regularity analysis near the interface, and the validation through a physically meaningful fourth-order example. These results provide a solid foundation for the numerical simulation of complex mechanical systems involving high-order differential operators and mixed boundary data.

References

1. Lions, J.L., Magenes, E. *Non-Homogeneous Boundary Value Problems and Applications*. Springer-Verlag, Berlin, 1972.
2. Grisvard, P. *Elliptic Problems in Nonsmooth Domains*. Society for Industrial and Applied Mathematics, Philadelphia, 2011.
3. Ciarlet, P.G. *The Finite Element Method for Elliptic Problems*. North-Holland, Amsterdam, 1978.
4. Brenner, S.C., Scott, L.R. *The Mathematical Theory of Finite Element Methods*. Springer, New York, 2008.
5. Dauge, M. *Elliptic Boundary Value Problems on Corner Domains*. Lecture Notes in Mathematics, Vol. 1341, Springer, Berlin, 1988.
6. Nazarov, S.A., Plamenevsky, B.A. *Elliptic Problems in Domains with Piecewise Smooth Boundaries*. Walter de Gruyter, Berlin, 1994.
7. RA Sobitov, JA Egamov. Some applications in economics. Web of Discoveries: Journal of Analysis and Inventions 2 (1), 32-35. January,2024
8. Sobitov R.A. Tanlanma va uning xarakteristikalari mavzusini o'qitishda pedagogik texnologiyalar tahlili. NamDU ilmiy axborotnomasi - Научный вестник НамГУ 2022 йил Maxsus сон. 692-694-бетлар
9. Sobitov R.A. Use Maple to teach Curves of the second order. International conference on contemporary mathematics and its applications - ICCMA 2021, p.102-103 pages.
10. Sobitov R.A Differential LG - game of many participant players**.** Middle European Scientific Bulletin. Volume - 16 (2021) September 125-132-pages.
11. BT Samatov, MA Xorilov, RA Sobitov, About one of the problem isaacs for lg– game Scientific Bulletin of Namangan State University 2 (11), 21-27

HSR (LONDON),
HOUGHTON STREET REVIEW JOURNAL
ISSN: 2058-5519|Impact factor-15.9
http://houghtonstreetreview.com/index.php/hsr

12. Sobitov R.A. Fur’e qatorlarining ba’zi tatbiqlari. NamDU ilmiy axborotnomasi - Научный вестник НамГУ 2024 йил 4-сон. 42-47-бетлар.
13. B.Irgashev, R.Sobitov. Riman-Liuvill ma’nosidagi hosila qatnashgan buzilishga ega oltinchi tartibli tenglama uchun aralash masala. Научный вестник НамГУ 2025 йил 12 son. 49-57-betlar